Research Signpost 37/661 (2), Fort P.O., Trivandrum-695 023, Kerala, India



Recent Res. Devel. Plant. Sci., 1(2003): 1-19 ISBN: 81-271-0047-1



The Xylariaceae as phytopathogens

Raymond L. Edwards¹, Nuchnart Jonglaekha², Anandini Kshirsagar³
Derek J. Maitland¹, Sureewan Mekkamol², Lianne K. Nugent⁴, Cherdchai Phosri⁴
Sureelak Rodtong⁵, Nutthaporn Ruchichachorn⁴, Ek Sangvichien⁶
George P. Sharples⁴, Prakitsin Sihanonth², Nuttika Suwannasai⁵
Surang Thienhirun⁶, Anthony J.S. Whalley⁴ and Margaret A. Whalley⁴
¹School of Chemistry, University of Bradford, Bradford, West Yorkshire, BD7 !DP, UK
²Royal Highland Project, 65 Doi Suthep Road, Chiang Mai, 50200, Thailand; ³MIT Project John Hopkins University, Pune 411001, India; ⁴School of Biomolecular Sciences, Liverpool John Moores University; Byrom Street, Liverpool L3 3AF, UK; ⁵Institute of Science Suranaree University of Technology, Nakhon Ratchasima, 30,000, Thailand; ⁴Faculty of Science, Ramkhamhaeng University, Bangkok, 10240, Thailand; ¹Department of Microbiology, Faculty of Science, Chulalongkorn University, Bangkok, 10300, Thailand
ðForest Products Research Division, Royal Forest Department, Chatuchak Bangkok 10900, Thailand

Abstract

This paper outlines the diseases caused by members of the Xylariaceae a large and well known ascomycete family with worldwide distribution. The significance of water stress in host trees in relation to development of canker disease following latent invasion is considered with examples from Europe, U.S.A, Australia, and Thailand. Root infections caused mainly by species of Rosellinia are reviewed and the implication of toxins produced by certain of these fungi is addressed. Reference to a third group of diseases, needle blights, is made.

Correspondence/Reprint request: Dr. Anthony J.S. Whalley, School of Biomolecular Sciences, Liverpool John Moores University; Byrom Street, Liverpool L3 3AF, UK. E-mail: A.J.WHALLEY@livjm.ac.uk