

# On discrete GB-splines

Boris I. Kvasov\*

(Received 7 August 2000)

## Abstract

Explicit formulae and recurrence relations are obtained for discrete generalized B-splines (discrete GB-splines for short). Properties of discrete GB-splines and their series are studied. It is shown that the series of discrete GB-splines is a variation diminishing function and the systems of discrete GB-splines are weak Chebyshev systems.

\* School of Mathematics, Suranaree University of Technology, Nakhon Ratchasima 30000, THAILAND. <mailto:boris@math.sut.ac.th>

<sup>0</sup>See <http://anziamj.austms.org.au/V42/CTAC99/Kvas> for this article and ancillary services, © Austral. Mathematical Soc. 2000. Published 27 Nov 2000.

Contents	C878
----------	------

## Contents

1 Introduction	C878
2 Discrete generalized splines	C879
3 Construction of discrete GB-splines	C882
4 Properties of discrete GB-splines	C885
5 Local approximation by discrete GB-splines	C887
6 Recurrence formulae for discrete GB-splines	C890
7 Series of discrete GB-splines (uniform case)	C892
References	C898

## 1 Introduction

The tools of generalized splines and GB-splines are widely used in solving problems of shape-preserving approximation (e.g., see [7]). Recently, in [1] a difference method for constructing shape-preserving hyperbolic tension splines as solutions of multipoint boundary value problems was developed. Such an approach permits us to avoid the computation of hyperbolic func-

tions and has substantial other advantages. However, the extension of a mesh solution will be a discrete hyperbolic tension spline.

The contents of this paper is as follows. In Section 2 we give a definition of a discrete generalized spline. Next, we construct a minimum length local support basis (whose elements are denoted as discrete GB-splines) of the new spline; see Section 3. Properties of GB-splines are discussed in Section 4, while the local approximation by discrete GB-splines of a given continuous function from its samples is considered in Section 5. In Section 6 we derive recurrence formulae for calculations with discrete GB-splines. The properties of GB-spline series are summarized in Section 7.

## 2 Discrete generalized splines

Let a partition  $\Delta : a = x_0 < x_1 < \dots < x_N = b$  of the interval  $[a, b]$  be given. We will denote by  $S_4^{DG}$  the space of continuous functions whose restriction to a subinterval  $[x_i, x_{i+1}]$ ,  $i = 0, \dots, N - 1$  is spanned by the system of four linearly independent functions  $\{1, x, \Phi_i, \Psi_i\}$ . In addition, we assume that each function in  $S_4^{DG}$  is smooth in the sense that for given  $\tau_i^{Lj} > 0$  and  $\tau_i^{Rj} > 0$ ,  $j = i - 1, i$ , the values of its first and second central divided differences with respect to the points  $x_i - \tau_i^{Lj}$ ,  $x_i$ ,  $x_i + \tau_i^{Rj}$  and  $x_i - \tau_i^{Lj}$ ,  $x_i$ ,  $x_i + \tau_i^{Rj}$  coincide.

Given a continuous function  $S$  we introduce the difference operators

$$\begin{aligned} D_1 S(x) \equiv D_{i,1} S(x) &= (\lambda_i^{Ri} S[x - \tau_i^{Li}, x] + \lambda_i^{Li} S[x, x + \tau_i^{Ri}]) (1 - t) \\ &\quad + (\lambda_{i+1}^{Ri} S[x - \tau_{i+1}^{Li}, x] + \lambda_{i+1}^{Li} S[x, x + \tau_{i+1}^{Ri}]) t, \\ D_2 S(x) \equiv D_{i,2} S(x) &= 2S[x - \tau_i^{Li}, x, x + \tau_i^{Ri}] (1 - t) \\ &\quad + 2S[x - \tau_{i+1}^{Li}, x, x + \tau_{i+1}^{Ri}] t, \\ &\quad x \in [x_i, x_{i+1}), \quad i = 0, \dots, N - 1, \end{aligned}$$

where  $\lambda_j^{Ri} = 1 - \lambda_j^{Li} = \tau_j^{Ri} / (\tau_j^{Li} + \tau_j^{Ri})$ ,  $j = i, i + 1$  and  $t = (x - x_i) / h_i$ ,  $h_i = x_{i+1} - x_i$ . The square parentheses denote the usual first and second divided differences of the function  $S$  with respect to the argument values  $x_j - \tau_j^{Li}$ ,  $x_j$ ,  $x_j + \tau_j^{Ri}$ ,  $j = i, i + 1$ .

**Definition 1** A discrete generalized spline is a function  $S \in S_4^{DG}$  such that

1. for any  $x \in [x_i, x_{i+1}]$ ,  $i = 0, \dots, N - 1$

$$\begin{aligned} S(x) \equiv S_i(x) &= [S(x_i) - \Phi_i(x_i) M_i] (1 - t) \\ &\quad + [S(x_{i+1}) - \Psi_i(x_{i+1}) M_{i+1}] t \\ &\quad + \Phi_i(x) M_i + \Psi_i(x) M_{i+1}, \end{aligned} \quad (1)$$

where  $M_j = D_{i,2} S_i(x_j)$ ,  $j = i, i + 1$ , and the functions  $\Phi_i$  and  $\Psi_i$  are subject to the constraints

$$\begin{aligned} \Phi_i(x_{i+1} - \tau_{i+1}^{Li}) &= \Phi_i(x_{i+1}) = \Phi_i(x_{i+1} + \tau_{i+1}^{Ri}) = 0, \\ \Psi_i(x_i - \tau_i^{Li}) &= \Psi_i(x_i) = \Psi_i(x_i + \tau_i^{Ri}) = 0, \\ D_{i,2} \Phi_i(x_i) &= 1, \quad D_{i,2} \Psi_i(x_{i+1}) = 1; \end{aligned} \quad (2)$$

2.  $S$  satisfies the following smoothness conditions

$$\begin{aligned} S_{i-1}(x_i) &= S_i(x_i), \\ D_{i-1,1}S_{i-1}(x_i) &= D_{i,1}S_i(x_i), \quad i = 1, \dots, N-1. \\ D_{i-1,2}S_{i-1}(x_i) &= D_{i,2}S_i(x_i), \end{aligned} \quad (3)$$

This definition generalizes the notion of a discrete polynomial spline in [9] and of a generalized spline in [5, 6]. The latter one can be obtained by setting  $\tau_j^{L_i} = \tau_j^{R_i} = 0$ ,  $j = i, i+1$  for all  $i$ . If  $\tau_i^{L_j} = \tau_i^L$  and  $\tau_i^{R_j} = \tau_i^R$ ,  $j = i-1, i$  then according to smoothness conditions (3) the values of the functions  $S_{i-1}$  and  $S_i$  at the three consecutive points  $x_i - \tau_i^L$ ,  $x_i$ ,  $x_i + \tau_i^R$  coincide. Setting  $\tau_j^{L_i} = \tau_j^{R_i} = \tau_i$ ,  $j = i, i+1$  we obtain  $D_{1,i}S(x) = S[x - \tau_i, x + \tau_i]$  and  $D_{2,i}S(x) = S[x - \tau_i, x, x + \tau_i]$ , which is the case discussed in [1].

The functions  $\Phi_i$  and  $\Psi_i$  depend on the tension parameters which influence the behaviour of  $S$  fundamentally. We call them the *defining functions*. In practice one takes  $\Phi_i(x) = \Phi_i(p_i, x)$ ,  $\Psi_i(x) = \Psi_i(q_i, x)$ ,  $0 \leq p_i, q_i < \infty$ . In the limiting case when  $p_i, q_i \rightarrow \infty$  we require that  $\lim_{p_i \rightarrow \infty} \Phi_i(p_i, x) = 0$ ,  $x \in (x_i, x_{i+1}]$  and  $\lim_{q_i \rightarrow \infty} \Psi_i(q_i, x) = 0$ ,  $x \in [x_i, x_{i+1})$  so that the function  $S$  in formula (1) turns into a linear function. Additionally, we require that if  $p_i = q_i = 0$  for all  $i$ , then we get a discrete cubic spline. If  $\tau_i^{L_j} = \tau_i^{R_j} = \tau_i$ ,  $j = i-1, i$  for all  $i$  then this spline coincides with a discrete cubic spline of [10]. The case  $\tau_i = \tau$  for all  $i$  was considered in [8].

### 3 Construction of discrete GB-splines

Let us construct a basis for the space of discrete generalized splines  $S_4^{DG}$  by using functions which have local supports of minimum length. Since  $\dim(S_4^{DG}) = 4N - 3(N-1) = N+3$  we extend the grid  $\Delta$  by adding the points  $x_j$ ,  $j = -3, -2, -1, N+1, N+2, N+3$ , such that  $x_{-3} < x_{-2} < x_{-1} < a$ ,  $b < x_{N+1} < x_{N+2} < x_{N+3}$ .

We demand that the discrete GB-splines  $B_i$ ,  $i = -3, \dots, N-1$  have the properties

$$\begin{aligned} B_i(x) &> 0, \quad x \in (x_i + \tau_i^{R_i}, x_{i+4} - \tau_{i+4}^{L_{i+3}}), \\ B_i(x) &\equiv 0, \quad x \notin (x_i, x_{i+4}), \end{aligned} \quad (4)$$

$$\sum_{j=-3}^{N-1} B_j(x) \equiv 1, \quad x \in [a, b]. \quad (5)$$

According to (1), on the interval  $[x_j, x_{j+1}]$ ,  $j = i, \dots, i+3$ , the discrete GB-spline  $B_i$  has the form

$$B_i(x) \equiv B_{i,j}(x) = P_{i,j}(x) + \Phi_j(x)M_{j,B_i} + \Psi_j(x)M_{j+1,B_i}, \quad (6)$$

where  $P_{i,j}$  is a polynomial of the first degree and  $M_{l,B_i} = D_{j,2}B_i(x_l)$ ,  $l = j, j+1$  are constants to be determined. The smoothness conditions (3) together with the constraints (2) give the following relations

$$\begin{aligned} P_{i,j}(x_j) &= P_{i,j-1}(x_j) + z_j M_{j,B_i}, \\ D_{j,1}P_{i,j}(x_j) &= D_{j-1,1}P_{i,j-1}(x_j) + c_{j-1,2}M_{j,B_i}, \end{aligned}$$

where

$$\begin{aligned} z_j &\equiv z_j(x_j) = \Psi_{j-1}(x_j) - \Phi_j(x_j), \\ c_{j-1,2} &= D_{j-1,1}\Psi_{j-1}(x_j) - D_{j,1}\Phi_j(x_j). \end{aligned}$$

Thus in (6)

$$P_{i,j}(x) = P_{i,j-1}(x) + [z_j + c_{j-1,2}(x - x_j)]M_{j,B_i}. \quad (7)$$

By repeated use of this formula we get

$$P_{i,j}(x) = \sum_{l=i+1}^j [z_l + c_{l-1,2}(x - x_l)]M_{l,B_i} = - \sum_{l=j+1}^{i+3} [z_l + c_{l-1,2}(x - x_l)]M_{l,B_i}.$$

As  $B_i$  vanishes outside the interval  $(x_i, x_{i+4})$ , we have from (7) that  $P_{i,j} \equiv 0$  for  $j = i, i + 3$ . In particular, the following identity is valid

$$\sum_{j=i+1}^{i+3} [z_j + c_{j-1,2}(x - x_j)]M_{j,B_i} \equiv 0,$$

from which one obtains the equalities

$$\sum_{j=i+1}^{i+3} c_{j-1,2} y_j^r M_{j,B_i} = 0, \quad r = 0, 1, \quad y_j = x_j - \frac{z_j}{c_{j-1,2}}. \quad (8)$$

Thus the formula for the discrete GB-spline  $B_i$  takes the form

$$B_i(x) = \begin{cases} \Psi_i(x)M_{i+1,B_i}, & x \in [x_i, x_{i+1}), \\ (x - y_{i+1})c_{i,2}M_{i+1,B_i} + \Phi_{i+1}(x)M_{i+1,B_i} \\ + \Psi_{i+1}(x)M_{i+2,B_i}, & x \in [x_{i+1}, x_{i+2}), \\ (y_{i+3} - x)c_{i+2,2}M_{i+3,B_i} + \Phi_{i+2}(x)M_{i+2,B_i} \\ + \Psi_{i+2}(x)M_{i+3,B_i}, & x \in [x_{i+2}, x_{i+3}), \\ \Phi_{i+3}(x)M_{i+3,B_i}, & x \in [x_{i+3}, x_{i+4}), \\ 0, & \text{otherwise.} \end{cases} \quad (9)$$

Substituting formula (9) into the normalization condition (5) written for  $x \in [x_i, x_{i+1}]$ , we obtain

$$\begin{aligned} \sum_{j=i-3}^i B_j(x) &= \Phi_i(x) \sum_{j=i-3}^{i-1} M_{i,B_j} + \Psi_i(x) \sum_{j=i-2}^i M_{i+1,B_j} \\ &+ (y_{i+1} - x)c_{i,2}M_{i+1,B_{i-2}} + (x - y_i)c_{i-1,2}M_{i,B_{i-1}} \equiv 1. \end{aligned}$$

As according to (5)

$$\sum_{j=i-3}^{i-1} M_{i,B_j} = \sum_{j=i-2}^i M_{i+1,B_j} = 0 \quad (10)$$

the following identity is valid

$$(y_{i+1} - x)c_{i,2}M_{i+1,B_{i-2}} + (x - y_i)c_{i-1,2}M_{i,B_{i-1}} \equiv 1.$$

From here one gets the equalities

$$y_{i+1}^r c_{i,2} M_{i+1, B_{i-2}} - y_i^r c_{i-1,2} M_{i, B_{i-1}} \equiv \delta_{1,r}, \quad r = 0, 1,$$

where  $\delta_{1,r}$  is the Kronecker symbol. Solving this system of equations and using (8) or (10), we obtain

$$\begin{aligned} M_{j, B_i} &= \frac{y_{i+3} - y_{i+1}}{c_{j-1,2} \omega'_{i+1}(y_j)}, \quad j = i+1, i+2, i+3, \\ \omega_{i+1}(x) &= (x - y_{i+1})(x - y_{i+2})(x - y_{i+3}) \end{aligned}$$

or with the notation  $c_{j,3} = y_{j+2} - y_{j+1}$ ,  $j = i, i+1$ ,

$$\begin{aligned} M_{i+1, B_i} &= \frac{1}{c_{i,2} c_{i,3}}, \\ M_{i+2, B_i} &= -\frac{1}{c_{i+1,2}} \left( \frac{1}{c_{i,3}} + \frac{1}{c_{i+1,3}} \right), \\ M_{i+3, B_i} &= \frac{1}{c_{i+2,2} c_{i+1,3}}. \end{aligned} \quad (11)$$

## 4 Properties of discrete GB-splines

The functions  $B_j$ ,  $j = -3, \dots, N-1$  possess many of the properties inherent in usual discrete polynomial B-splines. To provide inequality (4), in what follows we need to impose additional conditions on the functions  $\Phi_j$  and  $\Psi_j$ .

The proofs of the following four assertions repeat those given in [5].

**Lemma 2** *If the conditions*

$$\begin{aligned} 0 &< 2h_{j-1}^{-1} \Psi_{j-1}(x_j) < D_{j-1,1} \Psi_{j-1}(x_j), \\ 0 &< 2h_j^{-1} \Phi_j(x_j) < -D_{j,1} \Phi_j(x_j), \quad j = i+1, i+2, i+3 \end{aligned} \quad (12)$$

are satisfied, then in (11)  $c_{j,k} > 0$ ,  $j = i, \dots, i+4-k$ ;  $k = 2, 3$ , and

$$(-1)^{j-i-1} M_{j, B_i} > 0, \quad j = i+1, i+2, i+3. \quad (13)$$

**Theorem 3** *Let the conditions of Lemma 2 be satisfied, the functions  $\Phi_j$  and  $\Psi_j$  be convex and  $D_{j,2} \Phi_j$  and  $D_{j,2} \Psi_j$  be strictly monotone on the interval  $[x_j, x_{j+1}]$  for all  $j$ . Then the functions  $B_j$ ,  $j = -3, \dots, N-1$  have the following properties:*

1.  $B_j(x) > 0$  for  $x \in (x_j + \tau_j^{R_j}, x_{j+4} - \tau_{j+4}^{L_{j+3}})$ , and  $B_j(x) \equiv 0$  if  $x \notin (x_j, x_{j+4})$ ;
2.  $B_j$  satisfies the smoothness conditions (3);
3.  $\sum_{j=-3}^{N-1} y_{j+2}^r B_j(x) \equiv x^r$ ,  $r = 0, 1$  for  $x \in [a, b]$ ,  $\Phi_j(x) = c_{j-1,2} c_{j-2,3} B_{j-3}(x)$ ,  $\Psi_j(x) = c_{j,2} c_{j,3} B_j(x)$  for  $x \in [x_j, x_{j+1}]$ ,  $j = 0, \dots, N-1$ .

**Lemma 4** *The function  $B_i$  has support of minimum length.*

**Theorem 5** *The functions  $B_i$ ,  $i = -3, \dots, N - 1$ , are linearly independent and form a basis of the space  $S_4^{DG}$  of discrete generalized splines.*

## 5 Local approximation by discrete GB-splines

According to Theorem 5, any discrete generalized spline  $S \in S_4^{DG}$  can be uniquely written in the form

$$S(x) = \sum_{j=-3}^{N-1} b_j B_j(x) \quad (14)$$

for some constant coefficients  $b_j$ .

If the coefficients  $b_j$  in (14) are known, then by virtue of formula (9) we can write out an expression for the discrete generalized spline  $S$  on the interval  $[x_i, x_{i+1}]$ , which is convenient for calculations,

$$S(x) = b_{i-2} + b_{i-1}^{(1)}(x - y_i) + b_{i-1}^{(2)}\Phi_i(x) + b_i^{(2)}\Psi_i(x), \quad (15)$$

where

$$b_k^{(k)} = \frac{b_j^{(k-1)} - b_{j-1}^{(k-1)}}{c_{j,4-k}}, \quad k = 1, 2; \quad b_j^{(0)} = b_j. \quad (16)$$

The representations (14) and (15) allow us to find a simple and effective way to approximate a given continuous function  $f$  from its samples.

**Theorem 6** *Let a continuous function  $f$  be given by its samples  $f(y_j)$ ,  $j = -1, \dots, N + 1$ . Then for  $b_j = f(y_{j+2})$ ,  $j = -3, \dots, N - 1$ , formula (14) is exact for polynomials of the first degree and provides a formula for local approximation.*

**Proof:** It suffices to prove that the identities

$$\sum_{j=-3}^{N-1} y_{j+2}^r B_j(x) \equiv x^r, \quad r = 0, 1 \quad (17)$$

hold for  $x \in [a, b]$ . Using formula (15) with the coefficients  $b_{j-2} = 1$  and  $b_{j-2} = y_j$ ,  $j = i - 1, i, i + 1, i + 2$ , for an arbitrary interval  $[x_i, x_{i+1}]$ , we find that identities (17) hold.

For  $b_{j-2} = f(y_j)$ , formula (15) can be rewritten as

$$\begin{aligned} S(x) = & f(y_i) + f[y_i, y_{i+1}](x - y_i) + (y_{i+1} - y_{i-1})f[y_{i-1}, y_i, y_{i+1}]c_{i-1,2}^{-1}\Phi_i(x) \\ & + (y_{i+2} - y_i)f[y_i, y_{i+1}, y_{i+2}]c_{i,2}^{-1}\Psi_i(x), \quad x \in [x_i, x_{i+1}]. \end{aligned}$$

This is the formula of local approximation. The theorem is thus proved. ♠

**Corollary 7** *Let a continuous function  $f$  be given by its samples  $f_j = f(x_j)$ ,  $j = -2, \dots, N + 2$ . Then by setting*

$$b_{j-2} = f_j - \frac{1}{c_{j-1,2}} \left( \Psi_{j-1}(x_j) f[x_j, x_{j+1}] - \Phi_j(x_j) f[x_{j-1}, x_j] \right) \quad (18)$$

in (14), we obtain a formula of three-point local approximation, which is exact for polynomials of the first degree.

**Proof:** To prove the corollary, it is sufficient to take the monomials 1 and  $x$  as  $f$ . Then according to (18), we obtain  $b_{j-2} = 1$  and  $b_{j-2} = y_j$  and it only remains to make use of identities (17). This proves the corollary. ♠

Equation (15) permits us to write the coefficients of the spline  $S$  in its representation (14) of the form

$$b_{j-2} = \begin{cases} S(y_j) - D_{j-1,2} S(x_{j-1}) \Phi_{j-1}(y_j) - D_{j,2} S(x_j) \Psi_{j-1}(y_j), & y_j < x_j, \\ S(y_j) - D_{j,2} S(x_j) \Phi_j(y_j) - D_{j+1,2} S(x_{j+1}) \Psi_j(y_j), & y_j \geq x_j. \end{cases}$$

According to this formula we have  $b_{j-2} = S(y_j) + O(\bar{h}_j^2)$ ,  $\bar{h}_j = \max(h_{j-1}, h_j)$ . Hence it follows that the control polygon (e.g., see [4]) converges quadratically to the function  $f$  when  $b_{j-2} = f(y_j)$ , or if the formula (18) is used.

## 6 Recurrence formulae for discrete GB-splines

Let us define functions

$$B_{j,2}(x) = \begin{cases} D_{j,2} \Psi_j(x), & x \in [x_j, x_{j+1}), \\ D_{j+1,2} \Phi_{j+1}(x), & x \in [x_{j+1}, x_{j+2}], \\ 0, & \text{otherwise,} \end{cases} \quad j = i, i + 1, i + 2. \quad (19)$$

We assume that the functions  $D_{j,2} \Psi_j$  and  $D_{j+1,2} \Phi_{j+1}$  are strictly monotone on  $[x_j, x_{j+1})$  and  $[x_{j+1}, x_{j+2}]$  respectively. The splines  $B_{j,2}$  are a generalization of the ‘‘hat-functions’’ for polynomial B-splines. They are nonnegative and, furthermore,  $B_{j,2}(x_{j+l}) = \delta_{1,l}$ ,  $l = 0, 1, 2$ .

According to (9), (11) and (19) the function  $D_2 B_i$  can be written as

$$\begin{aligned} D_2 B_i(x) &= \sum_{j=i+1}^{i+3} M_{j,B_i} B_{j-1,2}(x) \\ &= \frac{1}{c_{i,3}} \left( \frac{B_{i,2}(x)}{c_{i,2}} - \frac{B_{i+1,2}(x)}{c_{i+1,2}} \right) - \frac{1}{c_{i+1,3}} \left( \frac{B_{i+1,2}(x)}{c_{i+1,2}} - \frac{B_{i+2,2}(x)}{c_{i+2,2}} \right). \end{aligned} \quad (20)$$

The function  $D_1 B_i$  satisfies the relation

$$D_1 B_i(x) = \frac{B_{i,3}(x)}{c_{i,3}} - \frac{B_{i+1,3}(x)}{c_{i+1,3}}, \quad (21)$$

where

$$B_{j,3}(x) = \begin{cases} \frac{D_{j,1}\Psi_j(x)}{c_{j,2}}, & x \in [x_j, x_{j+1}), \\ 1 + \frac{D_{j+1,1}\Phi_{j+1}(x)}{c_{j,2}} - \frac{D_{j+1,1}\Psi_{j+1}(x)}{c_{j+1,2}}, & x \in [x_{j+1}, x_{j+2}), \\ -\frac{D_{j+2,1}\Phi_{j+2}(x)}{c_{j+1,2}}, & x \in [x_{j+2}, x_{j+3}), \\ 0, & \text{otherwise.} \end{cases} \quad (22)$$

Using formula (22) it is easy to show that functions  $B_{j,3}$ ,  $j = -2, \dots, N-1$  satisfy the first and second smoothness conditions in (3), have supports of minimum length, are linearly independent and form a partition of unity,

$$\sum_{j=-2}^{N-1} B_{j,3}(x) \equiv 1, \quad x \in [a, b].$$

Applying formulae (20) and (21) to the representation (14) we also obtain

$$D_1 S(x) = \sum_{j=-2}^{N-1} b_j^{(1)} B_{j,3}(x), \quad D_2 S(x) = \sum_{j=-1}^{N-1} b_j^{(2)} B_{j,2}(x), \quad (23)$$

where  $b_j^{(k)}$ ,  $k = 1, 2$  are defined in (16).

## 7 Series of discrete GB-splines (uniform case)

Let us suppose that each step size  $h_i = x_{i+1} - x_i$  of the mesh  $\Delta : a = x_0 < x_1 < \dots < x_N = b$  is an integer multiple of the same tabulation step,  $\tau$ , of some detailed uniform refinement on  $[a, b]$ .

For  $\theta \in \mathbb{R}$ ,  $\tau > 0$  define

$$\mathbb{R}_{\theta\tau} = \{\theta + i\tau \mid i \text{ is an integer}\}$$

and let  $\mathbb{R}_{\theta 0} = \mathbb{R}$ . For any  $a, b \in \mathbb{R}$  and  $\tau > 0$  let

$$[a, b]_\tau = [a, b] \cap \mathbb{R}_{a\tau}.$$

The functions  $B_{j,2}$ ,  $B_{j,3}$ , and  $B_j$  with  $\tau_j^{Li} = \tau_j^{Ri} = \tau$ ,  $j = i, i+1$  for all  $i$  are nonnegative on the discrete interval  $[a, b]_\tau$ . This permits us to reprove the main results for discrete polynomial splines of [9] for series of discrete generalized splines. Even more, one can obtain the results of generalized splines of [5] from the corresponding statements for discrete generalized splines as a limiting case when  $\tau \rightarrow 0$ .

In particular, if in (14) and (23) we have the coefficients  $b_j^{(k)} > 0$ ,  $k = 0, 1, 2$ ,  $j = -3 + k, \dots, N-1$ , then the spline  $S$  will be a positive, monotonically increasing and convex function on  $[a, b]_\tau$ .

Let  $f$  be a function defined on the discrete set  $[a, b]_\tau$ . We say that  $f$  has

a zero at the point  $x \in [a, b]_\tau$  provided

$$f(x) = 0 \quad \text{or} \quad f(x - \tau) \cdot f(x) < 0.$$

When  $f$  vanishes at a set of consecutive points of  $[a, b]_\tau$ , say  $f$  is 0 at  $x, \dots, x + (r - 1)\tau$ , but  $f(x - \tau) \cdot f(x + r\tau) \neq 0$ , then we call the set  $X = \{x, x + \tau, \dots, x + (r - 1)\tau\}$  a *multiple zero* of  $f$ , and we define its multiplicity by

$$Z_X(f) = \begin{cases} r, & \text{if } f(x - \tau) \cdot f(x + r\tau) < 0 \text{ and } r \text{ is odd,} \\ r, & \text{if } f(x - \tau) \cdot f(x + r\tau) > 0 \text{ and } r \text{ is even,} \\ r + 1, & \text{otherwise.} \end{cases}$$

This definition assures that  $f$  changes sign at a zero if and only if the zero is of odd multiplicity.

Let  $Z_{[a, b]_\tau}(f)$  be the number of zeros of a function  $f$  on the discrete set  $[a, b]_\tau$ , counted according to their multiplicity. Let us denote  $D_1^L S(x) = S[x - \tau, x]$ .

**Theorem 8 (Rolle's Theorem For Discrete Generalized Splines.)** For any  $S \in S_4^{DG}$ ,

$$Z_{[a, b]_\tau}(D_1^L S) \geq Z_{[a, b]_\tau}(S) - 1. \quad (24)$$

**Proof:** First, if  $S$  has a  $z$ -tuple zero on the set  $X = \{x, \dots, x + (r - 1)\tau\}$ , it follows that  $D_1^L S$  has a  $(z - 1)$ -tuple zero on the set  $X' = \{x + \tau, \dots, x + (r - 1)\tau\}$ . Now if  $X^1$  and  $X^2$  are two consecutive zero sets of  $S$ , then it is trivially true that  $D_1^L S$  must have a sign change at some point between  $X^1$  and  $X^2$ . Counting all of these zeros, we arrive at the assertion (24). This completes the proof. ♠

**Lemma 9** Let the function  $D_{i,2}\Phi_i$  and  $D_{i,2}\Psi_i$  be strictly monotone on the interval  $[x_i, x_{i+1}]$  for all  $i$ . Then for every  $S \in S_4^{DG}$  which is not identically zero on any interval  $[x_i, x_{i+1}]_\tau$ ,  $i = 0, \dots, N - 1$ ,

$$Z_{[a, b]_\tau}(S) \leq N + 2.$$

**Proof:** According to (19) and (23), the function  $D_2 S$  has no more than one zero on  $[x_i, x_{i+1}]$ , because the functions  $D_2\Phi_i$  and  $D_2\Psi_i$  are strictly monotone and nonnegative on this interval. Hence  $Z_{[a, b]_\tau}(D_2 S) \leq N$ . Then according to the Rolle's Theorem 8, we find  $Z_{[a, b]_\tau}(S) \leq N + 2$ . This completes the proof. ♠

Denote by  $\text{supp}_\tau B_i = \{x \in \mathbb{R}_{a, \tau} \mid B_i(x) > 0\}$  the discrete support of the spline  $B_i$ , i.e. the discrete set  $(x_i + \tau, x_{i+4} - \tau)_\tau$ .

**Theorem 10** Assume that  $\zeta_{-3} < \zeta_{-2} < \dots < \zeta_{N-1}$  are prescribed points on the discrete line  $\mathbb{R}_{a, \tau}$ . Then

$$D = \det(B_i(\zeta_j)) \geq 0, \quad i, j = -3, \dots, N - 1$$

and strict positivity holds if and only if

$$\zeta_i \in \text{supp}_\tau B_i, \quad i = -3, \dots, N - 1. \quad (25)$$

The proof of this theorem is based on Lemma 9 and repeats that of Theorem 8.66 in [9, p.355]. The following statements follow immediately from Theorem 10.

**Corollary 11** *The system of discrete GB-splines  $\{B_j\}$ ,  $j = -3, \dots, N - 1$ , associated with knots on  $\mathbb{R}_{a,\tau}$  is a weak Chebyshev system according to the definition given in [9, p. 36], i.e. for any  $\zeta_{-3} < \zeta_{-2} < \dots < \zeta_{N-1}$  in  $\mathbb{R}_{a,\tau}$  we have  $D \geq 0$  and  $D > 0$  if and only if condition (25) is satisfied. In the latter case the discrete generalized spline  $S(x) = \sum_{j=-3}^{N-1} b_j B_j(x)$  has no more than  $N + 2$  zeros.*

**Corollary 12** *If the conditions of Theorem 5 are satisfied, then the solution of the interpolation problem*

$$S(\zeta_i) = f_i, \quad i = -3, \dots, N - 1, \quad f_i \in \mathbb{R} \quad (26)$$

*exists and is unique.*

Let  $A = \{a_{ij}\}$ ,  $i = 1, \dots, m$ ,  $j = 1, \dots, n$ , be a rectangular  $m \times n$  matrix with  $m \leq n$ . The matrix  $A$  is said to be totally nonnegative (totally

positive) (e.g., see [3]) if the minors of all order of the matrix are nonnegative (positive), i.e. for all  $1 \leq p \leq m$  we have

$$\det(a_{i_k j_l}) \geq 0 \quad (> 0) \quad \text{for all} \quad \begin{array}{l} 1 \leq i_1 < \dots < i_p \leq m, \\ 1 \leq j_1 < \dots < j_p \leq n. \end{array}$$

**Corollary 13** *For arbitrary integers  $-3 \leq \nu_{-3} < \dots < \nu_{p-4} \leq N - 1$  and  $\zeta_{-3} < \zeta_{-2} < \dots < \zeta_{p-4}$  in  $\mathbb{R}_{a,\tau}$  we have*

$$D_p = \det\{B_{\nu_i}(\zeta_j)\} \geq 0, \quad i, j = -3, \dots, p - 4$$

*and strict positivity holds if and only if*

$$\zeta_i \in \text{supp}_\tau B_{\nu_i}, \quad i = -3, \dots, p - 4$$

*i.e. the matrix  $\{B_j(\zeta_i)\}$ ,  $i, j = -3, \dots, N - 1$  is totally nonnegative.*

The last statement is proved by induction based on Theorem 5 and the recurrence relations for the minors of the matrix  $\{B_j(\zeta_i)\}$ . The proof does not differ from that of Theorem 8.67 described by [9, p.356].

Since the supports of discrete GB-splines are finite, the matrix of system (26) is banded and has seven nonzero diagonals in general. The matrix is tridiagonal if  $\zeta_i = x_{i+2}$ ,  $i = -3, \dots, N - 1$ .

An important particular case of the problem, in which  $S'(x_i) = f'_i$ ,  $i = 0, N$ , can be obtained by passing to the limit as  $\zeta_{-3} \rightarrow \zeta_{-2}$ ,  $\zeta_{N-1} \rightarrow \zeta_{N-2}$ .

De Boor and Pinkus [2] proved that linear systems with totally nonnegative matrices can be solved by Gaussian elimination without choosing a pivot element. Thus, the system (26) can be solved effectively by the conventional Gauss method.

Denote by  $S^-(\mathbf{v})$  the number of sign changes (variations) in the sequence of components of the vector  $\mathbf{v} = (v_1, \dots, v_n)$ , with zeros being neglected. Karlin [3] showed that if a matrix  $A$  is totally nonnegative then it decreases the variation, i.e.

$$S^-(A\mathbf{v}) \leq S^-(\mathbf{v}).$$

By virtue of Corollary 4, the totally nonnegative matrix  $\{B_j(\zeta_i)\}$ ,  $i, j = -3, \dots, N-1$ , formed by discrete GB-splines decreases the variation.

For a bounded real function  $f$ , let  $S^-(f)$  be the number of sign changes of the function  $f$  on the real axis  $\mathbb{R}$ , without taking into account the zeros

$$S^-(f) = \sup_n S^-[f(\zeta_1), \dots, f(\zeta_n)], \quad \zeta_1 < \zeta_2 < \dots < \zeta_n.$$

**Theorem 14** *The discrete generalized spline  $S(x) = \sum_{j=-3}^{N-1} b_j B_j(x)$  is a variation diminishing function, i.e. the number of sign changes of  $S$  does*

*not exceed that in the sequence of its coefficients:*

$$S^-\left(\sum_{j=-3}^{N-1} b_j B_j\right) \leq S^-(\mathbf{b}), \quad \mathbf{b} = (b_{-3}, \dots, b_{N-1}).$$

The proof of this statement does not differ from that of Theorem 8.68 for discrete polynomial B-splines in [9, p.356].

**Acknowledgements:** The author was supported by the Thailand Research Fund under grant number BRG/08/2543. He also wishes to express his gratitude to the referees for their invaluable remarks and suggestions for improvements.

## References

- [1] P. Costantini, B. I. Kvasov, and C. Manni. On discrete hyperbolic tension splines. *Advances in Computational Mathematics*, 11:331–354, 1999. C878, C881
- [2] C. De Boor and A. Pinkus. Backward error analysis for totally positive linear systems. *Numer. Math.*, 27:485–490, 1977. C897
- [3] S. Karlin. *Total Positivity, Volume 1*. Stanford University Press, Stanford, 1968. C896, C897

- [4] P. E. Koch and T. Lyche. Exponential B-splines in tension. In C.K. Chui, L.L. Schumaker, and J.D. Ward, editors, *Approximation Theory VI: Proceedings of the Sixth International Symposium on Approximation Theory. Vol. II*, pages 361–364, 1989. Academic Press. C889
- [5] B. I. Kvasov. Local bases for generalized cubic splines. *Russ. J. Numer. Anal. Math. Modelling*, 10:49–80, 1995. C881, C886, C892
- [6] B. I. Kvasov. GB-splines and their properties. *Annals of Numerical Mathematics*, 3:139–149, 1996. C881
- [7] B. I. Kvasov. Algorithms for shape preserving local approximation with automatic selection of tension parameters. *Computer Aided Geometric Design*, 17:17–37, 2000. C878
- [8] T. Lyche. *Discrete polynomial spline approximation methods*. PhD thesis, University of Texas, Austin, 1975. For a summary, see *Spline Functions*, K. Böhmer, G. Meinardus, and W. Schimpp, editors, Karlsruhe 1975, Lectures Notes in Mathematics No. 501, pages 144–176, Springer-Verlag, Berlin, 1976. C881
- [9] L. L. Schumaker. *Spline Functions: Basic Theory*. John Wiley & Sons, New York, 1981. C881, C892, C895, C895, C896, C898
- [10] Yu. S. Zav'yalov, B. I. Kvasov, and V. L. Miroshnichenko. *Methods of Spline Functions*. Nauka, Moscow (in Russian), 1980. C881