



# Relaxed Control for a Class of Strongly Nonlinear Impulsive Evolution Equations

PAIROTE SATTAYATHAM

School of Mathematics, Suranaree University of Technology  
Nakhon Ratchasima 30000, Thailand  
[sattayatham@yahoo.com](mailto:sattayatham@yahoo.com)

**Abstract**—Relaxed control for a class of strongly nonlinear impulsive evolution equations is investigated. Existence of solutions of strongly nonlinear impulsive evolution equations is proved and properties of original and relaxed trajectories are discussed. The existence of optimal relaxed control and relaxation results are also presented. For illustration, one example is given. © 2006 Elsevier Ltd. All rights reserved.

**Keywords**—Impulsive system, Banach space, Nonlinear monotone operator, Evolution triple, Relaxation.

## 1. INTRODUCTION

In this paper, we present sufficient conditions of optimality for optimal relaxed control problems arising in systems governed by strongly nonlinear impulsive evolution equations on Banach spaces. The general descriptions of such systems were proposed in [1] as given below.

$$\begin{aligned} \dot{x}(t) + A(t, x(t)) &= g(t, x(t), u(t)), & t \in I \setminus D, \\ x(0) &= x_0, \\ \Delta x(t_i) &= F_i(x(t_i)), & i = 1, 2, \dots, n, \end{aligned} \quad (1)$$

where  $I \equiv (0, T)$  is a bounded open interval of the real line and let the set  $D \equiv \{t_1, t_2, \dots, t_n\}$  be a partition on  $(0, T)$  such that  $0 < t_1 < t_2 < \dots < t_n < T$ . In general, the operator  $A$  is a nonlinear monotone operator,  $g$  is a nonlinear nonmonotone perturbation,  $\Delta x(t) \equiv x(t_i^+) - x(t_i^-) \equiv x(t_i^+) - x(t_i)$ ,  $i = 1, 2, \dots, n$ , and  $F_i$ s are nonlinear operators. This model includes all the standard models used by many authors in the field (see [2,3]). The objective functional is given by  $J(x, u) = \int_0^T L(t, x(t), u(t)) dt$ .

In a recent paper by the author [1], the existence of optimal control was proved, but sufficient conditions of relaxation for optimality were not addressed. We wish to present just that. Before we can consider such problems, we need some preparation. The rest of the paper is organized as follows. In Section 2, some basic notations and terminologies are presented. Section 3 contains

This work was supported by Thailand Research Fund. Grant No. BRG 48 2005.  
I would like to thank Prof. X.L. Xiang for her valuable suggestions and comments.

some preparatory results. Relaxed impulsive systems are presented in Section 4. Sufficient conditions of relaxation for optimality are discussed in Section 5. In Section 6, we presented an example for illustration.

## 2. SYSTEM DESCRIPTION

Let  $V$  be a real reflexive Banach space with topological dual  $V^*$  and  $H$  be a real separable Hilbert space. Let  $V \hookrightarrow H \hookrightarrow V^*$  be an evolution triple and the embedding  $V \hookrightarrow H$  be compact.

The system model considered here is based on this evolution triple (see [4, Chapter 23]).

Let  $\langle x, y \rangle$  denote the pairing of an element  $x \in V^*$  and an element  $y \in V$ . If  $x, y \in H$ , then  $\langle x, y \rangle = (x, y)$ , where  $(x, y)$  is the scalar product on  $H$ . The norm in any Banach space  $X$  will be denoted by  $\|\cdot\|_X$ .

Let  $p, q \geq 1$  be such that  $2 \leq p < +\infty$  and  $1/p + 1/q = 1$  and let  $I \equiv (0, T)$ . For  $p, q$  satisfying the preceding conditions, it follows from reflexivity of  $V$  that both  $L_p(I, V)$  and  $L_q(I, V^*)$  are reflexive Banach spaces. The pairing between  $L_p(I, V)$  and  $L_q(I, V^*)$  will be denoted by  $\langle \cdot, \cdot \rangle$ .

Define

$$W_{pq}(I) = \{x : x \in L_p(I, V), \dot{x} \in L_q(I, V^*)\},$$

with the norm

$$\|x\|_{W_{pq}(I)} = \|x\|_{L_p(L, V)} + \|\dot{x}\|_{L_q(L, V^*)},$$

where  $\dot{x}$  denoted the derivative of  $x$  in the generalized sense. The space  $(W_{pq}(I), \|\cdot\|_{W_{pq}(I)})$  becomes a Banach space which is clearly reflexive and separable and the embedding  $W_{pq}(I) \hookrightarrow C(\bar{I}, H)$  is continuous. If the embedding  $V \hookrightarrow H$  is compact, the embedding  $W_{pq}(I) \hookrightarrow L_p(I, H)$  is also compact (see [4, Problem 23.13(b)]). Sometimes we write  $W_{pq}(0, T)$  instead of  $W_{pq}(I)$ . Similarly, we can define  $W_{pq}(s, t)$  for  $0 \leq s < t \leq T$  and the space  $(W_{pq}(s, t), \|\cdot\|_{W_{pq}(s, t)})$  is also a separable reflexive Banach space. Moreover, the embedding  $W_{pq}(s, t) \hookrightarrow C([s, t], H)$  is continuous and the embedding  $W_{pq}(s, t) \hookrightarrow L_p((s, t), H)$  is also compact. We define the set  $PW_{pq}(0, T) = \{x : x|_{(t_i, t_{i+1})} \in W_{pq}(t_i, t_{i+1}); i = 0, 1, 2, \dots, n, t_0 = 0, t_{n+1} = T\}$ . For each  $x \in PW_{pq}(0, T)$ , we define  $\|x\|_{PW_{pq}(0, T)} = \sum_{i=0}^n \|x\|_{W_{pq}(t_i, t_{i+1})}$ . As a result, the space  $(PW_{pq}(0, T), \|\cdot\|_{PW_{pq}(0, T)})$  becomes a Banach space. Let  $PC([0, T], H) = \{x : x \text{ is a map from } [0, T] \text{ into } H \text{ such that } x \text{ is continuous at every point } t \neq t_i, \text{ left continuous at } t = t_i \text{ and right-hand limit } x(t_i^+) \text{ exists for } i = 1, 2, \dots, n\}$ . Equipped with the supremum norm topology, it is a Banach space. Consider the following impulsive evolution equation:

$$\begin{aligned} \dot{x}(t) + A(t, x(t)) &= g(t, x(t)), & t \in I \setminus D, \\ x(0) &= x_0 \in H, \\ \Delta x(t_i) &= F_i(x(t_i)), & i = 1, 2, \dots, n \text{ and } 0 < t_1 < t_2 < \dots < t_n < T. \end{aligned} \quad (2)$$

By a solution  $x$  of problem (2), we mean a function  $x \in PW_{pq}(0, T) \cap PC([0, T], H)$  such that  $x(0) = x_0$  and  $\Delta x(t_i) = F_i(x(t_i))$ ,  $i = 1, 2, \dots, n$ , and satisfies

$$\langle \dot{x}(t), v \rangle + \langle A(t, x(t)), v \rangle = \langle g(t, x(t)), v \rangle$$

for all  $v \in V$  and  $\mu$ -a.e. on  $I$ , where  $\mu$  is the Lebesgue measure on  $I$ .

## 3. PREPARATORY RESULTS

For the study of relaxation for optimality it is essential to guarantee the existence and uniqueness of solutions of the impulsive evolution equation and certain other related equations. Here in this section, for convenience of the reader, we quote some results from the recent work of the author [1]. First, we recall some hypotheses on the data of problem (2).

HYPOTHESIS (A).  $A : I \times V \rightarrow V^*$  is an operator such that

- (1)  $t \mapsto A(t, x)$  is weakly measurable, i.e., the functions  $t \mapsto \langle A(t, x), v \rangle$  are  $\mu$ -measurable on  $I$ , for all  $x, v \in V$ .
- (2) For each  $t \in I$ , the operator  $A(t) : V \rightarrow V^*$  is uniformly monotone and hemicontinuous, that is, there is a constant  $c_1 \geq 0$  such that

$$\langle A(t, x_1) - A(t, x_2), x_1 - x_2 \rangle \geq c_1 \|x_1 - x_2\|_V^p$$

for all  $x_1, x_2 \in V$ , and the map  $s \mapsto \langle A(t, x + sz), y \rangle$  is continuous on  $[0, 1]$  for all  $x, y, z \in V$ .

- (3) Growth condition: there exists a constant  $c_2 > 0$  and a nonnegative function  $a_1(\cdot) \in L_q(I)$  such that

$$\|A(t, x)\|_{V^*} \leq a_1(t) + c_2 \|x\|_V^{p-1},$$

for all  $x \in V$ , for all  $t \in I$ .

- (4) Coerciveness: there exists a constant  $c_3 > 0$  and  $c_4 \geq 0$  such that

$$\langle A(t, x), x \rangle \geq c_3 \|x\|_V^p - c_4, \quad \text{for all } x \in V, \text{ for all } t \in I.$$

HYPOTHESIS (G).  $g : I \times H \rightarrow V^*$  is an operator such that

- (1)  $g$  is measurable in the first variable and continuous in the second variable.
- (2)  $g$  is locally Lipschitz with respect to  $x$ , i.e., for any  $\rho > 0$ , there exists a positive constant  $L(\rho)$  such that

$$\|g(t, x_1) - g(t, x_2)\|_{V^*} \leq L(\rho) \|x_1 - x_2\|_H,$$

for all  $t \in I$  and all  $x_1, x_2 \in H$  with  $\|x_1\| \leq \rho, \|x_2\| \leq \rho$ .

- (3) There exist  $c_5 > 0, 2 \leq k < p$ , and a nonnegative function  $h_1(\cdot) \in L_q(I)$  such that

$$\|g(t, x)\|_{V^*} \leq h_1(t) + c_5 \|x\|_H^{k-1}$$

for all  $x \in V, t \in I$ .

HYPOTHESIS (F).  $F_i : H \rightarrow H$  is locally Lipschitz continuous on  $H$ , i.e., for any  $\rho > 0$ , there exists a constant  $L_i(\rho)$  such that

$$\|F_i(x_1) - F_i(x_2)\|_H \leq L_i(\rho) \|x_1 - x_2\|_H$$

for all  $x_1, x_2 \in H$  with  $\|x_1\|_H \leq \rho, \|x_2\|_H \leq \rho$  ( $i = 1, 2, \dots, n$ ).

It is sometimes convenient to rewrite system (2) into an operator equation. To do this, we set  $X = L_p(I, V)$  and hence  $X^* = L_q(I, V^*)$ . Moreover, we set

$$\begin{aligned} A(x)(t) &= A(t, x(t)), \\ G(x)(t) &= g(t, x(t)), \end{aligned} \tag{3}$$

for all  $x \in X$  and for all  $t \in (0, T)$ . Then the original equation (2) is equivalent to the following operator equation (see [4, Theorem 30.A]):

$$\begin{aligned} \dot{x} + Ax &= G(x), \\ x(0) &= x_0 \in H, \\ \Delta x(t_i) &= F_i(x(t_i)), \quad i = 1, 2, \dots, n \quad \text{and} \quad 0 < t_1 < t_2 < \dots < t_n < T. \end{aligned} \tag{4}$$

It follows from Theorem 30.A of Zeidler [4] that equation (4) defines an operator  $A : X \rightarrow X^*$  such that  $A$  is uniformly monotone, hemicontinuous, coercive, and bounded. Moreover, by using Hypothesis (G)(3) and using the same technique as in Theorem 30.A, one can show that the operator  $G : L_p(I, H) \rightarrow X^*$  is also bounded and satisfies

$$\|G(v)\|_{X^*} \leq M_1 + M_2 \|v\|_{L_p(I, H)}^{k-1}, \quad \text{for all } v \in L_p(I, H).$$

We state the following results from [1, Theorem B].

**THEOREM 3.1.** Under Assumptions (A), (F), and (G), system (2) has a unique solution  $x \in PW_{pq}(0, T) \cap PC([0, T], H)$  and there is a constant  $M > 0$  such that  $\|x\|_{PW_{pq}(0, T)} \leq M$  and  $\|x\|_{PC([0, T], H)} \leq M$ .

**PROOF.** See [1, Theorem B] for the proof of existence and boundedness. The uniqueness follows from Assumption (G)(2). To see this, suppose that system (2) has two solutions  $x_1, x_2$ . Then it follows from integration by part formula and monotonicity of  $A(t, x)$  that

$$\begin{aligned} \|x_1(t) - x_2(t)\|_H^2 &= 2 \int_0^t \langle \dot{x}_1(s) - \dot{x}_2(s), x_1(s) - x_2(s) \rangle_{V^* - V} ds \\ &= -2 \int_0^t \langle A(s, x_1(s)) - A(s, x_2(s)), x_1(s) - x_2(s) \rangle_{V^* - V} ds \\ &\quad + 2 \int_0^t \langle g(s, x_1(s)) - g(s, x_2(s)), x_1(s) - x_2(s) \rangle_{V^* - V} ds \\ &\leq 2 \int_0^t \langle g(s, x_1(s)) - g(s, x_2(s)), x_1(s) - x_2(s) \rangle_{V^* - V} ds \\ &\leq 2 \int_0^t \|g(s, x_1(s)) - g(s, x_2(s))\|_{V^*} \|x_1(s) - x_2(s)\|_V ds \\ &\leq 2L(\rho) \int_0^t \|x_1(s) - x_2(s)\|_H \|x_1(s) - x_2(s)\|_V ds \\ &\leq 2L(\rho)C_1 \int_0^t \|x_1(s) - x_2(s)\|_H^2 ds, \end{aligned}$$

for some positive constant  $C_1$ . By Gronwall's lemma, we get  $x_1(t) = x_2(t)$  for all  $t \in [0, T]$ . Hence  $x_1 = x_2$  and this prove the uniqueness of the solution of system (2).

Now, let us consider the corresponding control system. We model the control space by a separable complete metric space  $Z$  (i.e., a Polish space). By  $P_f$  ( $P_{fc}$ ), we denote a class of nonempty closed (closed and convex) subset of  $Z$ . Let  $I = (0, T)$ . Recall that a multifunction  $\Gamma : I \rightarrow P_f(Z)$  is said to be measurable if for each  $F \in P_f Z$ ,  $\Gamma^{-1}(F)$  is Lebesgue measurable in  $I$ . We defined  $S_\Gamma$  to be the set of all measurable selections of  $\Gamma(\cdot)$ , i.e.,

$$S_\Gamma = \{u : I \rightarrow Z \mid u(t) \text{ is measurable and } u(t) \in \Gamma(t), \mu\text{-a.e. } t \in I\},$$

where  $\mu$  is the Lebesgue measure on  $I$ . Note that the set  $S_\Gamma \neq \emptyset$  if  $\Gamma : I \rightarrow P_f(Z)$  is measurable (see [5, Theorem 2.23, p. 100]). Consider the following control systems:

$$\begin{aligned} \dot{x}(t) + A(t, x(t)) &= g(t, x(t), u(t)), \quad t \in I, \\ x(0) &= x_0 \in H, \\ \Delta x(t_i) &= F_i(x(t_i)), \quad i = 1, 2, \dots, n, \quad 0 < t_1 < t_2 < \dots < t_n < T. \end{aligned} \tag{5}$$

Here, we require the operator  $A$  and  $F_i$ s of (5) to satisfy Hypotheses (A), and (F), respectively, as in Section 3. We now give some new hypotheses for the remaining data.

**HYPOTHESIS (U).**  $U : I \rightarrow P_{fc}(Z)$  is a measurable multifunction satisfying  $U(\cdot) \subset F$ , where  $F$  is a compact subset of  $Z$ . For the admissible controls, we choose the set  $U_{ad} = S_U$ .

**HYPOTHESIS (G1).**  $g : I \times H \times Z \rightarrow V^*$  is an operator such that

- (1)  $t \mapsto g(t, x, z)$  is measurable, and the map  $(x, z) \mapsto g(t, x, z)$  is continuous on  $H \times Z$ .
- (2) For each fixed  $z$ ,  $g(t, x, z)$  is locally Lipschitz continuous with respect to  $x$  and uniformly in  $t$ .
- (3) There exist constants  $a, b > 0$  such that

$$\|g(t, x, z)\|_{V^*} \leq a + b\|x\|_H^{k-1},$$

for all  $x \in H$ ,  $t \in (0, T)$ , and  $z \in Z$ , where  $2 \leq k < p$ .

By Assumption (U), the control set  $S_U$  is nonempty and is called the class of original control. Now, let us define

- $X_0 = \{x \in PW_{pq}(I) \cap PC(\bar{I}, H) \mid x \text{ is a solution of (5) corresponding to } u\}$ .
- $X_0$  is called the class of original trajectories.
- $A_0 = \{(x, u) \in PW_{pq}(I) \cap PC(\bar{I}, H) \times S_U \mid x \text{ is a solution of (5) corresponding to } u\}$ .
- $A_0$  is called the class of admissible state-control pairs.

The following theorem guarantees that  $X_0 \neq \emptyset$ . Its proof follows immediately from Theorem 3.1 by defining the function  $g_u(t, x) = g(t, x, u)$  and noting that  $g_u$  satisfies all hypotheses of Theorem 3.1.

**THEOREM 3.2.** *Assume that Hypotheses (A),(F), (G1), and (U) hold. For every  $u \in S_U$ , equation (5) has a unique solution  $x(u) \in PW_{pq}(I) \cap PC(\bar{I}, H)$ . Moreover the set  $X_0$  is bounded in  $PW_{pq}(I) \cap PC(\bar{I}, H)$ , i.e.,  $\|x(u)\|_{PW_{pq}(0,T)} \leq M$  and  $\|x(u)\|_{PC([0,T],H)} \leq M$  for all  $u \in S_U$ .*

### 4. RELAXED IMPULSIVE SYSTEMS

We consider the following optimal control problem:

$$\inf \left\{ J(x, u) = \int_0^T L(t, x(t), u(t)) dt \right\} \tag{P}$$

subject to equation (5).

It is well known that, to solve the optimization problem involving (P) and obtain an optimal state-control pair, we need some kind of convexity hypothesis on the orientor field  $L(t, x(t), u(t))$ . If the convexity hypothesis is no longer satisfied, in order to get an optimal admissible pair, we need to pass to a larger system with measure control (or know as "relaxed control") in which the orientor field have been convexified. For this purpose, we introduce the relaxed control and the corresponding relaxed systems.

Let  $Z$  be a separable complete metric space (i.e., a Polish space) and  $B(Z)$  be its Borel  $\sigma$ -field. Let  $(\Omega, \Sigma, \mu)$  be a measure space. We will denote the space of probability measures on the measurable space  $(Z, B(Z))$  by  $M_+^1(Z)$ .

A Caratheodory integrand on  $\Omega \times Z$  is a function  $f : \Omega \times Z \rightarrow R$  such that  $f(\cdot, x)$  is  $\Sigma$ -measurable on  $\Omega$ ,  $f(\omega, \cdot)$  is continuous on  $Z$  for all  $\omega \in \Omega$ , and  $\sup\{|f(\omega, z)| : z \in Z\} \leq \alpha(\omega)$ , a.e., for some functions  $\alpha(\cdot) \in L_1(\Omega)$ . We denote the set of all Caratheodory integrands on  $\Omega \times Z$  by  $\text{Car}(\Omega, Z)$ . By a transition probability, we mean a function  $\lambda : \Omega \times B(Z) \rightarrow [0, 1]$  such that for every  $A \in B(Z)$ ,  $\lambda(\cdot, A)$  is  $\Sigma$ -measurable and for every  $\omega \in \Omega$ ,  $\lambda(\omega, \cdot) \in M_+^1(Z)$ . We use  $R(\Omega, Z)$  to denote the set of all transition probability from  $(\Omega, \Sigma)$  into  $(Z, B(Z))$ . Following Balder [6], we can define a topology on  $R(\Omega, Z)$  as follows: let  $f \in \text{Car}(\Omega, Z)$  and define

$$I_f(\lambda) = \int_{\Omega} \int_Z f(\omega, z) \lambda(\omega) (dz) d\mu(\omega). \tag{6}$$

The weak topology on  $R(\Omega, Z)$  is defined as the weakest topology for which all functionals  $I_f : R(\Omega, Z) \rightarrow R$ ,  $f \in \text{Car}(\Omega, Z)$ , are continuous.

Suppose  $\Omega = I = [0, T]$  and  $Z$  is a compact Polish space, then the space  $\text{Car}(I, Z)$  can be identified with the separable Banach space  $L_1(I, C(Z))$  where  $C(Z)$  is the space of all real-valued continuous functions on  $Z$ . To see this, we associate to each Caratheodory integrand  $\phi(\cdot, \cdot)$  the map  $t \mapsto \phi(t, \cdot) \in C(Z)$ . Let  $M(Z)$  be the space of all regular bounded countably additive measure defined on  $B(Z)$ . We note that  $M(Z)$  is a Banach space under the total variation norm, i.e.,  $\|\lambda\|_{M(Z)} = |\lambda|(Z)$ . Then by Riesz representation theorem, the dual  $[C(Z)]^*$  can be identified algebraically and metrically with  $M(Z)$ . The duality pair between  $M(Z)$  and  $C(Z)$  is given by

$$\langle \lambda, f \rangle = \int_Z f(z) \lambda(dz). \tag{7}$$

So  $M(Z)$  is a separable (see [7, p. 265]) dual Banach space and hence has a Radon-Nikodym property. This observation, combined with Theorem 1 of Diestel and Uhr [8, p. 98], tells us that

$$L_1(I, C(Z))^* = L_\infty(I, M(Z)).$$

So the weak topology on  $R(I, Z)$  coincides with the relative  $w^*(L_\infty(I, M(Z))), L_1(I, C(Z))$  topology. The duality pair between  $L_\infty(I, M(Z))$  and  $L_1(I, C(Z))$  is given by

$$\begin{aligned} \langle\langle \lambda, f \rangle\rangle &= \int_0^T \langle \lambda(t), f(t) \rangle dt \\ &= \int_0^T \int_Z f(t)(z) \lambda(t)(dz) dt \\ &= \int_0^T \int_Z f(t, z) \lambda(t)(dz) dt, \end{aligned} \tag{8}$$

which is the same formula as in (6) with  $f(t, z) \equiv f(t)(z)$ . This fact will be useful in the study of the relaxed control system where the control functions are transition probability.

Now we introduce some assumptions imposed on the class of relaxed control which will be denoted by  $S_\Sigma$ .

ASSUMPTION (U1).  $Z$  is a compact Polish space,  $U : I \rightarrow P_{fc}(Z)$  is a measurable multifunction.

Define  $\Sigma(t) = \{\lambda \in M_+^1(Z), \lambda(U(t)) = 1\}$  and let  $S_\Sigma \subseteq R(I, Z)$  be the set of transition probabilities on  $I \times B(Z)$  that are measurable selections of  $\Sigma(\cdot)$ . For any  $u \in S_U$ , we define the relaxation  $\delta_u \in S_\Sigma$  of  $u$  by  $\delta_u(t) \equiv$  Dirac probability measure at  $u(t)$ . Then we can identify  $S_U \subseteq S_\Sigma$ . From now on, we shall consider  $S_U$  and  $S_\Sigma$  as a subspace of the topological space  $R(I, Z)$  with the weak topology defined above.

We list two lemmas which will be useful in discussing relaxation problem. The proofs can be found in [7, Theorem IV 2.1] and [6, Corollary 3], respectively.

LEMMA 4.1. Suppose  $Z$  is a compact Polish space. Then  $S_\Sigma$  is convex, compact, and sequentially compact.

LEMMA 4.2.  $S_U$  is dense in  $S_\Sigma$ .

THEOREM 4.3. Let  $h : I \times H \times Z \rightarrow R$  be such that

1.  $t \mapsto (t, x, z)$  is measurable and  $(x, z) \mapsto h(t, x, z)$  is continuous.
  2.  $|h(t, x, z)| \leq \psi(t) \in L_1(I)$  for all  $(x, z) \in H \times Z$ .
- If  $x_k \rightarrow x \in C([0, T], H)$  then

$$\bar{h}_k(\cdot, \cdot) \rightarrow \bar{h}(\cdot, \cdot), \quad \text{in } L_1(I, C(Z)),$$

as  $k \rightarrow \infty$ , where  $\bar{h}_k(t, z) = h(t, x_k(t), z)$  and  $\bar{h}(t, z) = h(t, x(t), z)$ .

PROOF. The proof is similar to Lemma 3.3 in [3].

Next, let us consider this new larger system know as “relaxed impulsive system”

$$\begin{aligned} \dot{x}(t) + A(t, x(t)) &= \int_Z g(t, x(t), z) \lambda(t)(dz), & 0 \leq t \leq T, \\ x(0) &= x_0, \\ \Delta x(t_i) &= F_i(x(t_i)), & i = 1, 2, \dots, n. \end{aligned} \tag{9}$$

We will denote the set of trajectories of (9) by  $X_r$ , i.e.,  $X_r = \{x \in PW_{pq}(I) \cap PC(\bar{I}, H) \mid x \text{ is a solution of (9) corresponding to } \lambda \in S_\Sigma\}$ . Moreover, the set of admissible state-control pairs

of (9) will be denoted by  $A_r = \{(x, \lambda) \in PW_{pq}(I) \cap PC(\bar{I}, H) \times S_\Sigma \mid x \text{ is a solution of (9) corresponding to } \lambda \in S_\Sigma\}$ .

Note that  $X_0 \subseteq X_r$ , since  $S_U \subseteq S_\Sigma$  and if the hypotheses of Theorem 3.2 are satisfied,  $X_0 \neq \phi \Rightarrow X_r \neq \phi$ . To see this, given any relaxed control  $\lambda \in S_\Sigma$ , if we set  $\bar{g}(t, x(t), \lambda(t)) = \int_Z g(t, x(t), z)\lambda(t)(dz)$  then, working as in the proof of Theorem 3.2, one can show that there exists a relaxed admissible trajectory  $x(\lambda)$  corresponding to  $\lambda$ . We summarize the above discussion into a theorem.

**THEOREM 4.4.** *Assume that Hypotheses (A), (F), (G1), and (U1) hold. For every  $\lambda \in S_\Sigma$ , equation (9) has a unique solution  $x(\lambda) \in PW_{pq}(I) \cap PC(\bar{I}, H)$ . Moreover the set  $X_r$  is bounded in  $PW_{pq}(I) \cap PC(\bar{I}, H)$ , i.e.,  $\|x(\lambda)\|_{PW_{pq}(0,T)} \leq M$  and  $\|x(\lambda)\|_{PC([0,T],H)} \leq M$  for all  $\lambda \in S_\Sigma$ .*

The next theorem gives us a useful relation between  $X_0$  and  $X_r$ .

**THEOREM 4.5.** *If Assumptions (A), (F), (G1), and (U1) hold, then  $X_r = \bar{X}_0$  (closure is taken in  $PC(\bar{I}, H)$ ).*

Before proving this theorem, we need a lemma.

**LEMMA 4.6.** *If Assumptions (A), (F), (G1), and (U1) hold and  $\lambda_k \rightarrow \lambda$  in  $R(I, Z)$ , suppose that  $\{x_k, x\}$  is the solution of (9) corresponding to  $\{\lambda_k, \lambda\}$ , by working with a subsequence if necessary,  $x_k \rightarrow x$  in  $PC(\bar{I}, H)$  as  $k \rightarrow \infty$ .*

**PROOF.** Suppose that  $\lambda_k \rightarrow \lambda$  in  $R(I, Z)$  as  $k \rightarrow \infty$  and  $\{x_k, x\}$  is the solution of (9) corresponding to  $\{\lambda_k, \lambda\}$ . Since  $(x_k, \lambda_k) \in A_r$  for each positive integer  $k$ , then  $(x_k, u_k)$  must satisfy the equation

$$\begin{aligned} \dot{x}_k(t) + A(t, x_k(t)) &= \int_Z g(t, x_k(t), z)\lambda_k(t)(dz), \\ x_k(0) &= x_0 \in H, \\ \Delta x_k(t_i) &= F_i(x_k(t_i)), \quad i = 1, 2, \dots, n, \quad 0 < t_1 < t_2 < \dots < t_n < T. \end{aligned} \tag{10}$$

To finish the proof, we try to choose  $y \in X_r$  such that  $y$  is a solution of (9) corresponding to this  $\lambda$  and  $x_k \rightarrow y$  in  $PC(\bar{I}, H)$  as  $k \rightarrow \infty$ . The uniqueness property of the solution of (9) implies  $x = y$  and hence  $x_k \rightarrow x$  in  $PC(\bar{I}, H)$ . This proves that  $x \in \bar{X}_0$  and we are done. We shall do this by considering each case separately.

**CASE 1.** Find  $y$  on the interval  $(0, t_1)$ . For notational convenience, we let  $I_1 = (0, t_1)$ ,  $X_1 = L_p(I_1, V)$ , and  $X_1^* = L_q(I_1, V^*)$ . We note that  $X_1 = L_p(I_1, V)$  can be considered as a closed subspace of  $X = L_p(I, V)$ . Let  $x_k^1$  and  $\lambda_k^1$  be the restriction of the functions  $x_k, \lambda_k$  on the interval  $I_1$  ( $k = 1, 2, \dots$ ). Hence, by Theorem 4.4,  $\{x_k^1\}$  is bounded in  $W_{pq}(I_1)$ . By reflexivity of  $W_{pq}(I_1)$  there is a subsequence of  $\{x_k^1\}$ , again denoted by  $\{x_k^1\}$ , such that

$$x_k^1 \xrightarrow{w} x^1, \quad \text{in } W_{pq}(I_1), \quad \text{as } k \rightarrow \infty. \tag{11}$$

Since the embedding  $W_{pq}(I_1) \hookrightarrow X_1$  is continuous, the embedding  $W_{pq}(I_1) \hookrightarrow L_p(I_1, H)$  is compact and the operator  $A : X_1 \rightarrow X_1^*$  maps bounded sets to bounded sets, it follows from (11) that there exists a subsequence of  $\{x_k^1\}$ , again denoted by  $\{x_k^1\}$ , such that

$$\begin{aligned} x_k^1 &\xrightarrow{w} x^1 \quad \text{in } X_1, & \dot{x}_k^1 &\xrightarrow{w} \dot{x}^1, \quad \text{in } X_1^*, \\ Ax_k^1 &\xrightarrow{w} \xi, & & \text{in } X_1^*, \\ x_k^1 &\xrightarrow{s} x^1, & & \text{in } L_p(I_1, H), \end{aligned} \tag{12}$$

and, by [3, Theorem 3.B],  $x_k^1 \xrightarrow{s} x^1$  in  $C([0, t_1], H)$ , as  $k \rightarrow \infty$  and for some  $\xi \in X_1^*$ . Consider the following equation:

$$\begin{aligned} \dot{x}_k^1(t) + A(t, x_k^1(t)) &= \int_Z g(t, x_k^1(t), z)\lambda_k^1(t)(dz); & 0 \leq t < t_1, \\ x_k^1(0) &= x_0. \end{aligned} \tag{13}$$

Define an operator  $G_k : I_1 \rightarrow V^*$  and  $G : I_1 \rightarrow V^*$  as follows:

$$G_k(t) = \int_Z g(t, x_k^1(t), z) \lambda_k^1(t)(dz), \quad k = 1, 2, 3, \dots,$$

$$G(t) = \int_Z g(t, x^1(t), z) \lambda^1(t)(dz).$$

It follows from Assumption (G1) that  $G$  and  $G_k \in L_q(I_1, V^*)$ .

With this new notation, equation (13) can be rewritten into an equivalent operator equation of the form

$$\begin{aligned} \dot{x}_k^1 + A(x_k^1) &= G_k; & 0 < t < t_1, \\ x_k^1(0) &= x_0. \end{aligned} \tag{14}$$

For each fixed  $v \in V$ , define

$$\begin{aligned} \bar{g}_k(t, z) &= \langle g(t, x_k^1(t), z), v \rangle_{V^* - V}, \\ \bar{g}(t, z) &= \langle g(t, x^1(t), z), v \rangle_{V^* - V}. \end{aligned}$$

It follows from Assumption (G1) that, for each fixed  $t \in I_1$ ,

$$\bar{g}_k(t, \cdot), \text{ and } \bar{g}(t, \cdot) \in C(Z),$$

and furthermore

$$\bar{g}_k(\cdot, \cdot), \text{ and } \bar{g}(\cdot, \cdot) \in L_1(I_1, C(Z)).$$

Since  $x_k^1 \xrightarrow{s} x^1$  in  $C([0, t_1], H)$  (see equation (12)), then Theorem 4.3 gives

$$\bar{g}_k(\cdot, \cdot) \rightarrow \bar{g}(\cdot, \cdot), \quad \text{in } L_1(I_1, C(Z)), \quad \text{as } k \rightarrow \infty.$$

Since  $\lambda_k^1 \rightarrow \lambda^1$  in  $R(I_1, Z)$ , by equation (7), we have  $\lambda_k^1 \xrightarrow{w^*} \lambda^1$  in  $(L_1(I_1, C(Z)))^*$  as  $k \rightarrow \infty$ . Hence, it follows from Proposition 21.6(e) of Zeidler [4, p. 216] that

$$\langle\langle \lambda_k^1, \bar{g}_k \rangle\rangle \rightarrow \langle\langle \lambda^1, \bar{g} \rangle\rangle, \quad \text{as } k \rightarrow \infty.$$

This means that

$$\int_0^{t_1} \int_Z \langle g(t, x_k^1(t), z), v \rangle \lambda_k^1(t)(dz) dt \rightarrow \int_0^{t_1} \int_Z \langle g(t, x^1(t), z), v \rangle \lambda^1(t)(dz) dt \tag{15}$$

as  $k \rightarrow \infty$ . The convergence in (15) is true for all  $v \in V$  then we get

$$G_k \xrightarrow{w} G, \quad \text{as } k \rightarrow \infty, \quad \text{in } L_q(I_1, V^*).$$

By equation (12),  $x_k^1 \xrightarrow{s} x^1$  in  $C([0, t_1], H)$  and this implies  $x_k^1(0) \rightarrow x^1(0)$  in  $H$  as  $k \rightarrow \infty$ . Referring to the initial condition, we have  $x_k^1(0) = x_0 \in H$  for all  $k = 1, 2, 3, \dots$ . Thus,  $x^1(0) = x_0$ .

Up to this point, we can conclude that  $x^1$  satisfies the following equation:

$$\begin{aligned} \dot{x}^1(t) + \xi(t) &= \int_Z g(t, x^1(t), z) \lambda(t)(dz), \\ x^1(0) &= x_0 \in H. \end{aligned}$$

Next we aim to prove that  $\xi = Ax^1$  in  $X_1^*$ .



To prove this we note from equation (14) that

$$\begin{aligned} \langle\langle A(x_k^1), x_k^1 \rangle\rangle_{X_1^* - X_1} &= \langle\langle A(x_k^1), x^1 \rangle\rangle_{X_1^* - X_1} \\ &- \langle\langle \dot{x}_k^1, x_k^1 - x^1 \rangle\rangle_{X_1^* - X_1} + \langle\langle G_k, x_k^1 - x^1 \rangle\rangle_{X_1^* - X_1}. \end{aligned} \tag{16}$$

From integration by part formula, we have

$$\begin{aligned} \langle\langle \dot{x}_k^1, x_k^1 - x^1 \rangle\rangle_{X_1^* - X_1} &= \langle\langle \dot{x}^1, x_k^1 - x^1 \rangle\rangle_{X_1^* - X_1} \\ &+ \frac{1}{2} \left( \|x_k^1(t_1) - x^1(t_1)\|_H^2 - \|x_k^1(0) - x^1(0)\|_H^2 \right). \end{aligned} \tag{17}$$

Substitute (17) into (16) and noting that the second term on the right-hand side of (17) is always nonnegative, then we get

$$\begin{aligned} \langle\langle A(x_k^1), x_k^1 \rangle\rangle_{X_1^* - X_1} &\leq \langle\langle A(x_k^1), x^1 \rangle\rangle_{X_1^* - X_1} - \langle\langle \dot{x}^1, x_k^1 - x^1 \rangle\rangle_{X_1^* - X_1} \\ &+ \|x_k^1(0) - x^1(0)\|_H^2 + \langle\langle G_k, x_k^1 - x^1 \rangle\rangle_{X_1^* - X_1}. \end{aligned}$$

Therefore,

$$\lim_{k \rightarrow \infty} \langle\langle A(x_k^1), x_k^1 \rangle\rangle_{X_1^* - X_1} \leq \langle\langle \xi, x^1 \rangle\rangle_{X_1^* - X_1},$$

and hence  $A$  satisfies condition (M) (see [4, p. 474]). Then we have

$$A(x^1) = \xi.$$

Now we can say that  $x^1$  is the solution of the following equation:

$$\begin{aligned} \dot{x}^1(t) + A(t, x^1(t)) &= \int_Z g(t, x^1(t), z) \lambda(t)(dz), \\ x^1(0) &= x_0 \in H. \end{aligned}$$

This proves that  $x^1$  satisfies (9) on the interval  $(0, t_1)$  and  $x^1$  is the required  $y$  on  $(0, t_1)$ .

CASE 2. Find  $y$  on the interval  $(t_1, t_2)$ . The proof is similar to Case 1. Here, let  $I_2 = (t_1, t_2)$ ,  $X_2 = L_p(I, V)$ , and  $X_2^* = L_q(I, V^*)$ . Let  $x_k^2, u_k^2$  be the restriction of the functions  $x_k$  and  $u_k$  on the interval  $I_2$ , respectively ( $k = 1, 2, \dots$ ). It follows from equation (10) that  $(x_k^2, u_k^2)$  satisfies the operator equation

$$\begin{aligned} \dot{x}_k^2 + A(x_k^2) &= G_k; \quad t_1 < t < t_2, \\ x_k^2(t_1^+) &= x_k^2(t_1^-) + F_1(x_k^2(t_1)), \end{aligned} \tag{18}$$

where  $x_k^2(t_1^-) = x_k^2(t_1) = x_k^1(t_1)$  ( $k = 1, 2, 3, \dots$ ). By using the same proof as in Case 1, we get that

$$x_k^2 \xrightarrow{w} x^2, \quad \text{in } W_{pq}(t_1, t_2), \quad \text{and} \quad x_k^2 \xrightarrow{s} x^2, \quad \text{in } C([t_1, t_2], H), \quad \text{as } k \rightarrow \infty,$$

which implies that  $x_k^2(t_1^+) \rightarrow x^2(t_1^+)$  in  $H$  as  $k \rightarrow \infty$  and, moreover,  $x_2$  is also satisfied the operator equation

$$\dot{x}^2 + A(x^2) = G; \quad t_1 < t < t_2.$$

We are left to verify the initial condition at  $t_1$ . To see this, we note that the expression on the right-hand side of (18) converges to  $x^1(t_1) + F_1(x^1(t_1))$  as  $k \rightarrow \infty$  (see Hypothesis (F)). On the other hand, the left-hand side  $x_k^2(t_1^+) \rightarrow x^2(t_1^+)$  in  $H$  as  $k \rightarrow \infty$ . Hence,  $x^2(t_1^+) = x^1(t_1) + F_1(x^1(t_1)) \equiv x^2(t_1^-) + F_1(x^2(t_1))$ . This proves that  $x^2$  satisfies (9) on the interval  $(t_1, t_2)$  and  $x^2$  is the required  $y$  on  $(t_1, t_2)$ . Continuing this process we can find  $y$  on the interval  $(t_j, t_{j+1})$ ,  $j = 0, 1, \dots, n$ . By piecing them together from  $j = 1, 2, \dots, n$  and taking into account the impact

of jump, we obtain  $y$  which is the solution of (9) corresponding to the relaxed control  $\lambda$  satisfying  $x_k \rightarrow y$  in  $PC(\bar{I}, H)$  as  $k \rightarrow \infty$ . Since  $x = y$ ,  $x_k \rightarrow x$  in  $PC(\bar{I}, H)$  as  $k \rightarrow \infty$ . The proof of Lemma 4.6 is now complete.

PROOF OF THEOREM 4.5. First, we shall show that  $X_r \subseteq \bar{X}_0$ . Letting  $x \in X_r$ , then there exists  $\lambda \in S_\Sigma$  such that  $(x, \lambda) \in A_r$ . By virtue of density result Lemma 4.2, there exists a sequence  $\{u_k\} \in S_U$  such that  $\delta_{u_k} \rightarrow \lambda$  in  $R(I, Z)$ . Let  $x_k$  be the solution of (9) corresponding to  $u_k$ . Then we have a sequence  $\{(x_k, u_k)\} \subset A_0$ . Since  $(x_k, u_k) \in A_0$  for each positive integer  $k$ , then  $(x_k, u_k)$  must satisfy the equation

$$\begin{aligned} \dot{x}_k(t) + A(t, x_k(t)) &= \int_Z g(t, x_k(t), z) \delta_{u_k}(t)(dz), \\ x_k(0) &= x_0 \in H, \\ \Delta x_k(t_i) &= F_i(x_k(t_i)), \quad i = 1, 2, \dots, n, \quad k = 1, 2, 3, \dots, \\ 0 &< t_1 < t_2 < \dots < t_n < T. \end{aligned}$$

Applying Lemma 4.6, we get  $x_k \rightarrow x$  in  $PC(\bar{I}, H)$ . This proves that  $x \in \bar{X}_0$  and hence  $X_r \subseteq \bar{X}_0$ . Finally, we will show that  $X_r$  is closed in  $PC(\bar{I}, H)$ . Let  $\{x_k\}$  be a sequence of points in  $X_r$  such that  $x_k \rightarrow x$  in  $PC(\bar{I}, H)$  as  $k \rightarrow \infty$ . By definition of  $X_r$ , there is a sequence  $\{\lambda_k\}$  of points in  $S_\Sigma$  such that  $(x_k, \lambda_k) \in A_r$ ,  $k = 1, 2, 3, \dots$ . By Lemma 4.1,  $S_\Sigma$  is compact in  $R(I, Z)$  under the weak topology. Moreover,  $R(I, Z)$ -topology coincides with the relative  $w^*(L_\infty(I, M(Z)), L_1(I, C(Z)))$  topology which is metrizable (see [2, p. 276]). Then, by passing to a subsequence if necessary, we may assume that  $\lambda_k \rightarrow \lambda$  in  $R(I, Z)$ . Applying Lemma 4.6, there is  $x \in X_r$  such that  $x_k \rightarrow x$  in  $PC(\bar{I}, H)$  as  $k \rightarrow \infty$ . Hence  $X_r$  is closed in  $PC(\bar{I}, H)$  and, consequently,  $\bar{X}_0 \subseteq \bar{X}_r = X_r$ . The proof of Theorem 4.5 is now complete.

The following corollary is an immediate consequence of Lemma 4.6.

COROLLARY 4.7. Under assumption of Theorem 4.5, the function  $\lambda \mapsto x(\lambda)$  is continuous from  $S_\Sigma \subseteq R(I, Z)$  into  $PC(\bar{I}, H)$ .

### 5. EXISTENCE OF OPTIMAL CONTROLS

Consider the following Lagrange optimal control problem (P<sub>r</sub>): find a control policy  $\bar{\lambda} \in S_\Sigma$ , such that it imparts a minimum to the cost functional  $J$  given by

$$J(\lambda) \equiv J(x^\lambda, \lambda) \equiv \int_I \int_Z l(t, x^\lambda(t), z) \lambda(t)(dz) dt, \tag{P<sub>r</sub>}$$

where  $x^\lambda$  is the solution of system (9) corresponding to the control  $\lambda \in S_\Sigma$ . We make the following hypothesis concerning the integrand  $l(\cdot, \cdot, \cdot)$ .

HYPOTHESIS (L).  $l : I \times H \times Z \rightarrow R \cup \{+\infty\}$  is Borel measurable satisfying the following conditions:

- (1)  $(\xi, z) \mapsto l(t, \xi, z)$  is lower semicontinuous on  $H \times Z$  for each fixed  $t$ .
- (2)  $\psi(t) \leq l(t, \xi, z)$  almost everywhere with  $\psi(t) \in L_1(I)$ .

Let  $m_r = \inf\{J(\lambda) : \lambda \in S_\Sigma\}$ . We have the following theorem on the existence of optimal impulsive control.

THEOREM 5.1. Suppose Assumptions (A), (F), (G1), (U1), (L) hold and  $Z$  is compact Polish space, then there exists  $(\bar{x}, \bar{\lambda}) \in A_r$  such that  $J(\bar{x}, \bar{\lambda}) = m_r$ .

PROOF. If  $J(\lambda) = +\infty$  for all  $\lambda \in S_\Sigma$ , then every control is admissible. Assume  $\inf\{J(\lambda) : \lambda \in S_\Sigma\} = m_r < +\infty$ . By Assumption (L), we have  $m_r > -\infty$ . Hence  $m_r$  is finite. Let  $\{\lambda_k\}$  be a minimizing sequence so that  $\lim_{k \rightarrow \infty} J(\lambda_k) = m_r$ . By Lemma 4.1,  $S_\Sigma$  is compact in the topology

$R(I, Z)$ . Hence, by passing to a subsequence if necessary, we may assume that  $\lambda_k \rightarrow \bar{\lambda}$  in  $R(I, Z)$  as  $k \rightarrow \infty$ . This means that  $\lambda_k \xrightarrow{w^*} \bar{\lambda}$  in  $L_\infty(I, M(Z))$  as  $k \rightarrow \infty$ . Let  $\{x_k, \bar{x}\}$  be the solution of (9) corresponding to  $\{\lambda_k, \bar{\lambda}\}$ . By Lemma 4.6, we get  $x_k \rightarrow \bar{x}$  in  $PC(I, H)$  and  $(\bar{x}, \bar{\lambda}) \in A_r$ . Next, we shall prove that  $(\bar{x}, \bar{\lambda})$  is an optimal pair.

As before, we identify the space of Caratheodory integrand  $\text{Car}(I, Z)$  with the separable Banach space  $L_1(I, C(Z))$ . We note that every semicontinuous measurable integrand  $l : I \times H \times Z \rightarrow R \cup \{+\infty\}$  is the limit of an increasing sequence of Caratheodory integrand  $\{l_j\} \in L_1(I, C(Z))$  for each fixed  $h \in H$ . Thus, there exists an increasing sequence of Caratheodory integrand  $\{l_j\} \in L_1(I, C(Z))$  such that

$$l_j(t, \bar{x}(t), z) \uparrow l(t, \bar{x}(t), z), \quad \text{as } j \rightarrow \infty, \quad \text{for all } t \in I, \quad z \in Z.$$

Since  $x_k \rightarrow \bar{x}$  in  $PC(I, H)$ , by applying Theorem 4.3 on each subinterval of  $[0, T]$ ,  $l_j(t, x_k(t), z) \rightarrow l_j(t, \bar{x}(t), z)$  as  $k \rightarrow \infty$  for almost all  $t \in I$  and all  $z \in Z$ . We note that since  $\lambda_k \xrightarrow{w^*} \bar{\lambda}$  in  $L_\infty(I, M(Z))$  as  $k \rightarrow \infty$ , then

$$\begin{aligned} J(\bar{x}, \bar{\lambda}) &= \langle \langle \bar{\lambda}, l \rangle \rangle = \int_I \int_Z l(t, \bar{x}(t), z) \bar{\lambda}(t)(dz) dt \\ &= \lim_{j \rightarrow \infty} \int_I \int_Z l_j(t, \bar{x}(t), z) \bar{\lambda}(t)(dz) dt \\ &= \lim_{j \rightarrow \infty} \lim_{k \rightarrow \infty} \int_I \int_Z l_j(t, x_k(t), z) \lambda_k(t)(dz) dt \\ &\leq \lim_{k \rightarrow \infty} \lim_{j \rightarrow \infty} \int_I \int_Z l_j(t, x_k(t), z) \lambda_k(t)(dz) dt = m_r. \end{aligned}$$

However, by definition of  $m_r$ , it is obvious that  $J(\bar{x}, \bar{\lambda}) \geq m_r$ . Hence  $J(\bar{x}, \bar{\lambda}) = m_r$ . This implies that  $(\bar{x}, \bar{\lambda})$  is an optimal pair.

REMARK. If  $J_0(x, u) = \int_I l(t, x(t), u(t)) dt$  is the cost functional for the original problem and  $m = \inf\{J_0(x, u) : u \in U_{\text{ad}}\}$ , in general we have  $m_r \leq m$ . It is desirable that  $m_r = m$ , i.e., our relaxation is reasonable. With some stronger conditions on  $l$ , i.e., the map  $(\xi, \eta, z) \mapsto l(t, \xi, z)$  is continuous and  $|l(t, \xi, z)| \leq \theta_R(t)$  for all most all  $t \in I$  and  $\theta_R \in L_1(I)$ , one can show that  $m_r = m$ . The proof is similar to Theorem 4.B. in [3].

### 6. EXAMPLE

In this section we present an example of a strongly nonlinear impulsive system for which our general theory can be applied. Let  $I = (0, T)$  and  $\Omega \subset \mathbb{R}^N$  be a bounded domain with  $C^1$  boundary  $\partial\Omega$ . For  $p \geq 2$  and  $\theta \geq 0$ , we consider the following quasi-linear parabolic control problem:

$$\begin{aligned} \frac{\partial}{\partial t} x(t, z) - \sum_{i=1}^N D_i \left( |D_i x(t, z)|^{p-2} D_i x(t, z) \right) &= g(t, z, x(t, z), u), & \text{on } I \times \Omega, \\ x|_{I \times \partial\Omega} &= 0, & x(0, z) = x_0(z), \\ \Delta x(t_i, z) &= F_i(x(t_i, z)), & i = 1, 2, \dots, n, \end{aligned} \tag{19}$$

where  $(0 < t_1 < t_2 < \dots < t_n < T)$ .

Here the operator  $D_i = \frac{\partial}{\partial x_i}$  ( $i = 1, 2, \dots, N$ ). We need the following hypotheses on the data of (19).

HYPOTHESIS (G').  $f : I \times \Omega \times R \times R^N \rightarrow R$  is an operator such that

- (1)  $(t, z) \mapsto f(t, z, \xi, u)$  is measurable  $I \times \Omega$ ; and the map  $(\xi, u) \mapsto f(t, z, \xi, u)$  is continuous on  $R \times R^N$ .

(2) There exist constants  $b_1 > 0$ ,  $b_2(t, z) \in L_2(I \times \Omega)$  such that

$$|f(t, z, \xi, u)| \leq b_1|\xi| + b_2(t, z).$$

HYPOTHESIS (F').  $F_i : L_2(\Omega) \rightarrow L_2(\Omega)$  ( $i = 1, 2, \dots, n$ ) are operators such that for any  $\rho > 0$  there exists a constant  $L_i(\rho) > 0$  such that

$$\|F_i(h_1) - F_i(h_2)\|_{L_2(\Omega)} \leq L_i(\rho) \|h_1 - h_2\|_{L_2(\Omega)}$$

for all  $h_1, h_2 \in L_2(\Omega)$  with  $\|h_1\|_{L_2(\Omega)} \leq \rho$ ,  $\|h_2\|_{L_2(\Omega)} \leq \rho$  ( $i = 1, 2, \dots, n$ ).

THEOREM 6.1. If Hypotheses (G') and (F') hold and letting  $x_0(\cdot) \in L_2(\Omega)$ ,  $u \in U_{\text{ad}}$  (defined below), then system (19) has a solution  $x \in L_p(I, PW^{1,p}(\Omega) \cap PC(\bar{I}, L_2(\Omega)))$  such that  $\frac{\partial x}{\partial t} \in L_q(I, W^{-1,q}(\Omega))$ .

PROOF. In this system, the evolution triple is  $V = W_0^{1,p}(\Omega)$ ,  $H = L_2(\Omega)$ , and  $V^* = W^{-1,q}(\Omega)$ . All embeddings are compact (Sobolev embedding theorem). Define an operator  $A : I \times V \rightarrow V^*$  by

$$\langle A(t, x), y \rangle_V = \int_{\Omega} \sum_{i=1}^N |D_i(x)|^{p-2} (D_i x)(D_i y) dz. \quad (20)$$

One can easily check that  $A(t, x)$  satisfies Hypothesis (A) in Section 3. The uniform monotonicity of  $A(t, \cdot)$  is a consequence of the result of Zeidler [4, p. 783].

In the sequel, we suppose that  $\beta_{1i}(\cdot)$ ,  $\beta_{2i}(\cdot)$  ( $1 \leq i \leq N$ ) are continuous functions from  $[0, T]$  to  $R$  and satisfy  $\beta_{1i}(\cdot) \leq \beta_{2i}(\cdot)$  ( $1 \leq i \leq N$ ) for all  $t \in [0, T]$ ,  $1 \leq i \leq N$ . There exists a constant  $a > 0$  such that  $-a \leq \beta_{1i}(t) \leq \beta_{2i}(t) \leq a$  for all  $t \in [0, T]$ ,  $1 \leq i \leq N$ . Set  $Z = [-a, a]^N \subset R^N$ . Then  $Z$  is a compact Polish space. Define

$$U(t) = \{(w_i(t)) \in R^N : \beta_{1i}(t) \leq w_i(t) \leq \beta_{2i}(t), 1 \leq i \leq N\}.$$

It is clear that  $U : I \rightarrow P_{fc}(Z)$  is measurable. The set of admissible controls  $U_{\text{ad}}$  is chosen as  $U_{\text{ad}} \equiv S_U \equiv \{u : I \rightarrow R^N \mid u \text{ is measurable and } u(t) \in U(t) \text{ a.e. } t \in [0, T]\}$ . Hence the multifunction  $U$  satisfies (U1).

Next, for  $t \in I$ ,  $\phi \in H$ ,  $w \in Z$  define a function  $b^w : I \times H \times V \rightarrow R$  by  $b^w(t, \phi, \psi) = \int_{\Omega} f(t, z, \phi, w)\psi(z) dz$ . Then, the map  $\psi \mapsto b^w(t, \phi, \psi)$  is bounded on  $V$  and hence is a continuous linear form on  $V$ . Thus there exists an operator  $g : I \times H \times Z \rightarrow V^*$  such that

$$b^w(t, \phi, \psi) = \langle g(t, \phi, w), \psi \rangle_{V^* - V}. \quad (21)$$

By using Hypothesis (G'), we obtain that  $g$  satisfies Hypothesis (G1) of Section 3.

Using the operator  $A$  and  $g$  as defined in equation (20) and (21), one can rewrite system (19) in an abstract form as in (9). So, applying Theorem 4.4, system (19) has a solution.

## REFERENCES

1. P. Sattayatham, Strongly nonlinear impulsive evolution equations and optimal control, *Nonlinear Analysis* **52**, 1005–1020, (2004).
2. N.S. Papageorgiou, Properties of the relaxed trajectories of evolution equations and optimal control, *SIAM J. Control Optim.* **27**, 267–288, (1989).
3. X. Xiang, P. Sattayatham and W. Wei, Relaxed controls for a class of strongly nonlinear delay evolution equations, *Nonlinear Analysis* **52**, 703–723, (2003).
4. E. Zeidler, *Nonlinear Functional Analysis and Its Applications*, Vol. I, II, Springer, New York, (1990).
5. X. Li and J. Yong, *Optimal Control Theory for Infinite Dimensional Systems*, Birkhäuser, Boston, MA, (1995).
6. E. Balder, A general denseness result for relaxed control theory, *Bull. Austral. Math. Soc.* **20**, 463–475, (1984).
7. J. Warga, *Optimal Control of Differential and Functional Equations*, Academic Press, New York, (1972).
8. J. Diestel and J.J. Uhr, *Vector Measures*, American Mathematical Society, (1997).