

## โครงการวิจัย: การศึกษาโครงสร้างและและสมบัติในการใช้ไคตินของเอนไซม์ไคติเนส เอ กลายพันธุ์ จากเชื้อแบคทีเรีย *Vibrio carchariae*

### บทคัดย่อ

เอนไซม์ไคติเนส เอ จากเชื้อแบคทีเรียในทะเล *Vibrio harveyi* เป็นเอนไซม์ที่สลายไคตินให้เป็นน้ำตาลไคโตโอลิโกแซคคารีไรต์และไคโตไบโอสเป็นผลิตภัณฑ์หลัก งานวิจัยก่อนหน้าได้ทำการโคลนยีนไคติเนส เอ และได้ศึกษาคุณสมบัติทางเอนไซม์ไคติเนส เอ งานวิจัยนี้อธิบายบทบาทของกรดอะมิโนวงแหวนที่บริเวณจับกับสับสเตรทและกรดอะมิโนที่ผิวของเอนไซม์ในการสลายสับสเตรทไคตินและไคโตโอลิโกแซคคารีไรต์โดยทำการเปลี่ยนกรดอะมิโน Trp70 Trp168 Tyr171 Trp231 Tyr245 Trp275 Trp397 และ Trp570 การทดสอบหาค่า specific hydrolyzing activity ของไคติเนสกลายพันธุ์ พบว่ามีเอนไซม์กลายพันธุ์ตัวเดียวคือ W397F ที่ให้ค่าแอกติวิตี้สูงกว่าเอนไซม์ดั้งเดิมส่วนเอนไซม์กลายพันธุ์อื่นมีค่าแอกติวิตี้ลดลงอย่างมาก การวิเคราะห์น้ำตาลผลิตภัณฑ์ที่สร้างขึ้นโดยวิธี TLC พบว่าเมื่อกรดอะมิโนที่ตำแหน่งรีดิวซ์ Trp275 ถูกเปลี่ยนเป็น Gly และ Trp397 เปลี่ยนเป็น Phe ทำให้รูปแบบการสลายน้ำตาลสายสั้นเปลี่ยนไปอย่างสิ้นเชิงแสดงว่ากรดอะมิโนทั้งสองน่าจะมี ความสำคัญต่อการเลือกจับของน้ำตาลไคโตโอลิโกแซคคารีไรต์ การศึกษาการจับกับไคตินและการทดลองทางจลนพลศาสตร์แสดงให้เห็นว่า Trp70 ซึ่งพบอยู่ที่ผิวที่ปลายด้านเอ็นของโดเมนจับไคตินมีความสำคัญมากต่อการจับกับไคตินสายยาว การตรวจหารูปแบบการจับของสับสเตรทโดยเทคนิค HPLC MS พบว่า NAG<sub>6</sub> ชอบบริเวณจับ -2 ถึง +2 มากกว่าบริเวณจับ -3 ถึง +2 ส่วน NAG<sub>5</sub> จะจับกับบริเวณจับ -2 ถึง +2 อย่างเดียว ในขณะที่ crystalline  $\alpha$  chitin จะเริ่มจับที่บริเวณจับได้หลายตำแหน่งทำให้สลายตัวกลางไคโตโอลิโกแซคคารีไรต์ ได้หลายชนิดซึ่งตัวกลางเหล่านี้จะจับกับตำแหน่ง -2 ถึง +2 เป็นหลัก นอกจากนี้ยังพบว่าไคติเนสกลายพันธุ์ W275G และ W397F มีความชอบต่อสับสเตรทชนิด  $\beta$  มากกว่าสับสเตรทชนิด  $\alpha$

## ABSTRACT

Chitinase A from a marine bacterium *Vibrio harveyi* is an enzyme that degrades chitin to chitooligosaccharides, yielding chitobiose as the major product. The gene encodes chitinase A was previously cloned and its enzymatic properties characterized. This study describes the functional roles of the aromatic residues located at the substrate binding cleft and the surface-exposed residues in chitin and chitooligosaccharide hydrolyses. Point mutations of Trp70, Trp168, Tyr171, Trp231, Tyr245, Trp275, Trp397, and Trp570, were generated. Investigation of specific hydrolyzing activity indicated that only mutant W397F had enhanced activity, while other mutants showed a significant loss of the activity. TLC analysis of product formation showed a complete change in the hydrolytic patterns of short-chain substrates when the reducing end residues Trp275 was mutated Gly and Trp397 to Phe, suggesting that both residues were crucial for the binding selectivity of chitinoligosaccharides. Chitin binding assay and kinetic experiments suggested that Trp70, which is located on the surface at the *N*-terminal end of the chitin binding domain, was the essential binding residue for a long-chain chitin. Assessment of substrate binding modes by HPLC MS revealed that NAG<sub>6</sub> preferred subsites -2 to +2 over subsites -3 to +2 and NAG<sub>5</sub> only bound to subsites -2 to +2. Crystalline  $\alpha$  chitin initially occupied various subsites, generating various chitooligosaccharide intermediates which later interacted mainly to subsites -2 to +2. In addition, mutants W275G and W397F preferred  $\beta$  substrates over  $\alpha$  substrates.