

ACKNOWLEDGEMENT

The completion of this work could not have been possible without the financial support of the National Research Council of Thailand (NRCT), the Royal Golden Jubilee Ph.D. (RGJ-PHD) Program (Grant No. NRCT5-RGJ63007-093, Suranaree University of Technology (SUT), Thailand Science Research and Innovation (TSRI), and National Science, Research and Innovation Fund (NSRF). The author also deeply appreciates the Center of Excellence on Technology and Innovation for Korat Chicken Business Development for their invaluable support.

I would like to express my deepest gratitude to my advisor, Asst. Prof. Dr. Wittawat Molee, for his invaluable mentorship, unwavering support, insightful suggestions, and continuous encouragement throughout my study. His guidance and wisdom have been instrumental in the successful completion of this work.

I extend my heartfelt thanks to my co-advisors, Assoc. Prof. Dr. Anyanee Kamkaew, School of Chemistry, Institute of Science, Assoc. Prof. Dr. Amonrat Molee, School of Animal Technology and Innovation, Institute of Agricultural Technology, Suranaree University of Technology, and Dr. Elisabeth Baéza, French National Institute for Agriculture, Food, and Environment (INRAE), UMR Biologie des Oiseaux et Aviculture, France for their expert advice, kind assistance, and ongoing support.

Special appreciation goes to my thesis committee members, Assoc. Prof. Dr. Amonrat Molee, Assoc. Prof. Dr. Anyanee Kamkaew, Assoc. Prof. Dr. Sutisa Khempaka, Asst. Prof. Dr. Pakanit Kupittayanant, Dr. Elisabeth Baéza and Dr. Cécile Berri, for their constructive feedback and valuable comments, which have significantly enhanced the quality and scientific rigor of this thesis.

I would like to express my gratitude to Dr. Teerapong Yata, Biochemistry Unit, Department of Physiology, Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand, for his guidance and support in providing lipid-based nanoparticles for the experiments in this study. Special thanks are also extended to Dr. Worapapar Treesuppharat and Apipu Ariyachayut, Thammasat University Research Unit in Mechanisms of Drug Action and Molecular Imaging, Drug Discovery and

Development Center, Thammasat University, Pathumthani, Thailand, for their valuable advice and assistance with imaging using the In vivo imaging system.

I am deeply grateful to Suranaree University of Technology (SUT) for providing essential resources, including the poultry unit, SUT farm, nutrition laboratory, and the Center for Scientific and Technological Equipment.

I would also like to thank my undergraduate and PhD colleagues, Wunpen Sonsamrong, Nitipol Chainet, Wichuta Khosinklang, Thanidtha Morawong, Sirirrot Khotsalee, and whose assistance with data collection and analysis was invaluable to the success of this study, for their collaboration, friendship, and encouragement throughout this journey.

Finally, I am forever grateful to my family for their unwavering love, support, and belief in me. Their encouragement has been my greatest source of strength. Lastly, I extend my sincere gratitude to everyone who supported me throughout this journey, even if their names do not appear here.

Piyaradtana Homyok